

# Markov Network Independencies


Sargur Srihari  
srihari@cedar.buffalo.edu

# Topics

- Markov Network Independencies
  - Basic Independencies
  - Independencies Revisited
  - From Distributions to Graphs

# Markov network captures independencies of interactions

Misconception MN with factors



$$P(a,b,c,d) = \frac{1}{Z} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

where

$$Z = \sum_{a,b,c,d} \phi_1(a,b) \cdot \phi_2(b,c) \cdot \phi_3(c,d) \cdot \phi_4(d,a)$$

Factorization of Distribution and Independences

$P$  supports  $(X \perp Y|Z)$  iff we can write distribution as  $P(X) = \Phi_1(X,Z) \Phi_2(Y,Z)$

$$P(A,B,C,D) = \left[ \frac{1}{Z} \phi_1(A,B) \cdot \phi_2(B,C) \right] \phi_3(C,D) \cdot \phi_4(D,A)$$

Factor with  $\{B, \{A, C\}\}$


Factor with  $\{D, \{A, C\}\}$

Therefore  $(B \perp D|A, C)$

Similarly  $(A \perp C|B, D)$

# Basic Independencies

- As in Bayesian Networks, graph structure in a Markov network encodes a set of independence assumptions
- In a MN Probabilistic influence flows along the undirected paths in the graph and “blocked” if we condition on intervening nodes
  - i.e., we know their values
- We state this formally, next




# Active Path Definition

- Let  $H$  be a Markov network with nodes  $\chi = \{X_1, \dots, X_n\}$ ,
- Let  $X_1 - \dots - X_k$  be a path in  $H$
- Let  $Z \subseteq \chi$  be a set of observed variables
- A Path  $X_1 - \dots - X_k$  is active given  $Z$  if none of the  $X_i$  is in  $Z$

– Example:

- Consider path  $A-D-C$
  - If  $Z = \{B\}$ , path  $A-D-C$  is active
- We can define separation in the graph when there is no active path, next




# Separation and Global Independencies

- Set of nodes  $Z$  separates sets  $X$  and  $Y$  denoted  $sep_H(X;Y|Z)$  if there is no active path between any  $X \in X$  and  $Y \in Y$ 
  - Example:
  - $Z = \{B, D\}$  separates  $A$  and  $C$ 
    - i.e.,  $sep_H(A;C|B,D)$  or there is no active path between  $A$  and  $C$

## Global independencies associated with $H$ are

- $I(H) = \{(X \perp Y | Z) : sep_H(X;Y|Z)\}$
- Independencies in  $I(H)$  guaranteed to hold for every distribution  $P$  over  $H$



$$I(H) = \{(A \perp C | B, D), (B \perp D | A, C)\}$$

# Definition of Separation Leads to a disadvantage

- With a superset of  $Z$ , separation still holds
  - If  $sep_H(X; Y|Z)$  then  $sep_H(X; Y|Z')$  for any  $Z' \supset Z$
- If separation is taken as definition of independencies, we restrict ability to encode non-monotonic independence relations
  - Non-monotonic reasoning is quite useful
    - E.g., intercausal reasoning with BNs
      - Two diseases are independent, but dependent given some common symptom
  - Such independence properties cannot be expressed as a Markov network

# Factorization and Independencies

- Can show connection between independence properties implied by a Markov structure and factorizing a distribution over the graph
- Analogous to Bayesian Networks
  - Let  $G$  be a BN for a set of random variables  $\chi$  and  $P$  be a distribution over  $\chi$ .
  - If  $P$  factorizes according to  $G$ , i.e., product of CPDs, then  $G$  is an I-map of  $P$ 
    - i.e., independencies  $I(G) \subseteq I$



# Formalizing independencies in MNs and distributions

- Gibbs Distribution

- A distribution  $P_\Phi$  is a Gibbs distribution parameterized by a set of factors  $\Phi = \{\phi_1(D_1), \dots, \phi_K(D_K)\}$
- If defined as follows

$$P_\Phi(X_1, \dots, X_n) = \frac{1}{Z} \tilde{P}(X_1, \dots, X_n)$$

where

$$\tilde{P}(X_1, \dots, X_n) = \prod_{i=1}^m \phi_i(D_i)$$

is an unnormalized measure and

$$Z = \sum_{X_1, \dots, X_n} \tilde{P}(X_1, \dots, X_n)$$
 is a normalizing constant

called the partition function

$D_i$  are sets of random variables

# Soundness of Separation Criterion

- Theorem 1 (factorization to independencies):
  - Let  $P$  be a distribution over  $\chi = \{X_1, \dots, X_n\}$  and  $H$  is a Markov structure over  $\chi$
  - If  $P$  is a Gibbs distribution that factorizes over  $H$ , then  $H$  is an I-map for  $P$ 
    - Factorize means every  $D_i$  in  $H$  is a clique
- Theorem 2: Hammersley-Clifford  
(other direction: independencies to factorization)
  - if  $H$  is an I-map for  $P$  then  $P$  factorizes over  $H$
  - Holds only for positive distributions ( $P > 0$  for every assignment)

# Positive Distribution


- A distribution  $P$  is said to be positive if for all events  $\alpha \in S$  such that  $\alpha \neq \Phi$  we have that  $P(\alpha) > 0$
- A non-positive distribution
- 16 possible values
  - Distribution  $P$ : 8 have value 1/8 rest are zero

$X_1$	$X_2$	$X_3$	$X_4$	$P(X)$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

# Non-positive distribution consistent with H

- Four binary random variables
- Global Independencies in graph H:
  - Consider  $X_1 - X_2 - X_3 - X_4 - X_1$ 
    - implies  $(X_1 \perp X_3 | X_2, X_4)$
- $P$  also satisfies this
  - For the assignment  $X_2=1, X_4=0$
  - $P(X_1=1 | X_2=1, X_4=0) = 1$ 
    - Rest are zero
    - Thus  $X_1$  is independent of  $X_3$




$X_1$	$X_2$	$X_3$	$X_4$	$P(\mathbf{X})$
0	0	0	0	1/8
1	0	0	0	1/8
1	1	0	0	1/8
1	1	1	0	1/8
0	0	0	1	1/8
0	0	1	1	1/8
0	1	1	1	1/8
1	1	1	1	1/8

Rest 8 probs are 0

- Global independencies hold  $\Rightarrow$  H is an I-map for P
- But  $P$  does not factorize according to H (Proof by contradiction)<sup>12</sup>

# Graphical Model as Filter



$p(x)$  is allowed to pass through only if  
It satisfies independencies in graph  
This set is denoted  $DF$  or  $UF$  (for BN or MN)

- $UI$  is set of distributions that are consistent with set of conditional independence statements read from the graph using graph separation
- $UF$  are set of distributions that can be expressed as factorization of the form

$$p(\mathbf{x}) = \frac{1}{Z} \prod_c \psi_c(\mathbf{x}_c)$$

- *Hammersley-Clifford theorem* states that  $UI$  and  $UF$  are identical

# Independencies in Bayesian Networks

- Bayesian networks have two types of independencies
  - Local independencies
    - Each node is independent of its non-descendants given its parents
  - Global independencies
    - Induced by d-separation
- These two sets of independencies are equivalent
  - One implies the other

# Three Independencies of Markov Networks

- Two local independencies
  - Pairwise  $I_p(H)$
  - Markov Blanket  $I_l(H)$
- Global independency  $I(H)$ 
  - Identify three sets of nodes  $A$ ,  $B$  and  $C$
  - To test conditional independence property
$$A \perp B | C$$
  - Consider all possible paths from nodes in set  $A$  to nodes in set  $B$ 
    - If all such paths pass through one or more nodes in  $C$  then path is blocked and independence holds


# Pairwise Independency $I_p(H)$

- Weakest independency
  - Whenever two variables are directly connected
    - Potential of being correlated not mediated by other variables
  - When two variables are not directly linked
    - There must be a way of rendering them conditionally independent
- If  $H$  is a Markov network its *pairwise independencies* are

$$I_p(H) = \{(X \perp Y | \chi - \{X, Y\}) : X - Y \notin H\}$$

- In the example

$$I_p(H) = \{(A \perp C | B, D), (B \perp D | A, C)\}$$




When  $X, Y$  are not directly connected they are independent given all other variables



# Markov Blanket Independency $I_l(H)$

- Analogous to local independencies in Bayesian networks
  - We can block all influences by conditioning on its immediate neighbors
    - Node is conditionally independent of all nodes given its immediate neighbors
- For graph  $H$  the Markov blanket of  $X$  in  $H$  is the set of neighbors of  $X$  in  $H$
- Local independencies associated are

$$I_l(H) = \{(X \perp \chi - \{X\} - MB_H(X) \mid MB_H(X)) : X \in \chi\}$$



# Relationship between Markov properties

- Three independencies of network structure  $H$
- $I_p(H)$  is strictly weaker than  $I_l(H)$  is strictly weaker than  $I(H)$
- For positive distributions all three are equivalent

# Separation in Markov Networks

- Markov network encodes a set of conditional independencies

- Probabilistic influence flows

- in undirected paths

- Blocked if we condition on intervening nodes

- Every path from any node in  $A$  to  $B$  passes through  $C$

- No explaining away


- Testing for independence simpler than in directed graphs

- Alternative view


- Remove all nodes in set  $C$  together with all their connecting links

- If no paths from  $A$  to  $B$  then conditional independence holds

- Markov blanket



- A node is conditionally independent of all other nodes conditioned only on its neighbors



# Factorization Properties

- Factorization rule corresponds to conditional independence test
- Notion of locality needed
- Consider two nodes  $x_i$  and  $x_j$  not connected by a link
  - They are conditionally independent given all other nodes in graph
    - Because there is no direct path between them and
    - All other paths pass through nodes that are observed and hence those paths are blocked
  - Expressed as
 

$$p(x_i, x_j \mid \mathbf{X} \setminus \{i, j\}) = p(x_i \mid \mathbf{X} \setminus \{i, j\}) p(x_j \mid \mathbf{X} \setminus \{i, j\})$$
  - Where  $\mathbf{X} \setminus \{i, j\}$  denotes set  $\mathbf{x}$  of all variables with  $x_i$  and  $x_j$  removed
- For conditional independence to hold
  - factorization is such that  $x_i$  and  $x_j$  do not appear in the same factor
    - No path between them other than going through others
  - leads to graph concept of clique

