

Bayes Decision Theory

Sargur Srihari

CSE 555
Introduction to Pattern Recognition

Reverend Thomas Bayes

1702-1761




Bayes set out his theory of probability in *Essay towards solving a problem in the doctrine of chances* published in the *Philosophical Transactions of the Royal Society of London* in 1764. The paper was sent to the Royal Society by Richard Price, a friend of Bayes', who wrote:-
I now send you an essay which I have found among the papers of our deceased friend Mr Bayes, and which, in my opinion, has great merit... In an introduction which he has writ to this Essay, he says, that his design at first in thinking on the subject of it was, to find out a method by which we might judge concerning the probability that an event has to happen, in given circumstances, upon supposition that we know nothing concerning it but that, under the same circumstances, it has happened a certain number of times, and failed a certain other number of times.

Bayes Rule

Two Classes(A, \sim A) , Single Binary-Valued Feature (X, \sim X)

Known

Prior probability:	$p(A) : 33.0\%$
Conditional probabilities:	
$p(X A) : 75.0\%$	
$p(X \sim A) : 25.0\%$	
Posterior probability:	
$p(A X) : 59.6\%$	
Visualization	: frequency
Result	: X
Reset	



Sample:	1000
A:	330
\sim A:	670
A & X:	248
A & \sim X:	82
\sim A & X:	168
\sim A & \sim X:	502

Data

By Conditional Probability Rule,

$$p(X / A) = \frac{p(X \ & \ A)}{p(A)} = \frac{.248}{.330} = 0.7515$$

$$p(X / \sim A) = \frac{p(X \ & \ \sim A)}{p(\sim A)} = \frac{.168}{.670} = 0.2507$$

By Bayes Rule, $P(A / X) = \frac{P(X / A)P(A)}{P(X)}$

$$\begin{aligned} &= \frac{P(X / A)P(A)}{P(X \ & \ A) + P(X \ & \ \sim A)} \\ &= \frac{P(X / A)P(A)}{P(X / A)P(A) + P(X / \sim A)P(\sim A)} \\ &= \frac{0.75 \times 0.33}{0.75 \times 0.33 + 0.25 \times 0.67} \\ &= \frac{.2475}{.2475 + .1675} = \frac{.2475}{.415} = 0.596 \end{aligned} \quad 2$$

Bayes Decision Theory

- Fundamental statistical approach to statistical pattern classification
- Quantifies trade-offs between classification using probabilities and costs of decisions
- Assumes all relevant probabilities are known

Prior Probabilities

State of nature, prior

- State of nature is a random variable


$$P(\omega_1) + P(\omega_2) = 1 \text{ (exclusivity and exhaustivity)}$$

- Decision rule with only the prior information

Decide ω_1 if $P(\omega_1) > P(\omega_2)$ otherwise decide ω_2

Class-conditional Probabilities

$$p(x / \omega_1) \text{ and } p(x / \omega_2)$$



Pdfs show the probability of measuring a particular feature value given category ω_j .

If x is a feature value, the two curves describe the difference in populations of two types of classes.

Density functions are normalized-- thus area under each curve is 1.0

Bayes formula to combine prior and class-conditional probabilities

$$P(\omega_j | x) = \frac{p(x | \omega_j)P(\omega_j)}{p(x)}$$


- In the case of two categories

$$p(x) = \sum_{j=1}^{j=2} P(x | \omega_j)P(\omega_j)$$

- Informally, Bayes rule says:
posterior = likelihood x prior / evidence


Posterior probabilities

Class-conditional
p.d.f.s



Feature x

Posterior probabilities for the priors $P(\omega_1) = 2/3, P(\omega_2) = 1/3$
For $x = 14$,
 $P(\omega_1/x) = 0.08, P(\omega_2/x) = 0.92$



Feature x

Bayes Decision Rule

x is an observation for which:

if $P(\omega_1 | x) > P(\omega_2 | x)$  True state of nature = ω_1

if $P(\omega_1 | x) < P(\omega_2 | x)$  True state of nature = ω_2

Therefore:

whenever we observe a particular x , the probability of error is :

$P(\text{error} | x) = P(\omega_1 | x) \text{ if we decide } \omega_2$

$P(\text{error} | x) = P(\omega_2 | x) \text{ if we decide } \omega_1$

Bayes Decision Rule minimizes probability of error

Decide ω_1 if $P(\omega_1 | x) > P(\omega_2 | x)$;
otherwise decide ω_2

Therefore:

$$P(\text{error} | x) = \min [P(\omega_1 | x), P(\omega_2 | x)]$$

(Bayes decision)

Bayes Decision Theory – Continuous Features

- Generalization of the preceding ideas
 - Use of more than one feature
 - Use more than two states of nature
 - Allowing actions and not only decide on the state of nature
 - Introduce a loss of function which is more general than the probability of error

Loss Function

- Allowing actions other than classification primarily allows the possibility of rejection
- Refusing to make a decision in close or bad cases!
- The loss function states how costly each action taken is

Loss Function Definition

Let $\{\omega_1, \omega_2, \dots, \omega_c\}$ be the set of c states of nature (or “categories”)

Let $\{\alpha_1, \alpha_2, \dots, \alpha_a\}$ be the set of possible actions

Let $\lambda(\alpha_i | \omega_j)$ be the loss incurred for taking action α_i when the state of nature is ω_j

Overall Risk

$R = \text{Sum of all } R(\alpha_i | x) \text{ for } i = 1, \dots, a$

Conditional risk

Minimizing $R \longleftrightarrow$ Minimizing $R(\alpha_i | x)$ for $i = 1, \dots, a$

Expected Loss with action i

$$R(\alpha_i | x) = \sum_{j=1}^{j=c} \lambda(\alpha_i | \omega_j) P(\omega_j | x)$$

Select the action α_i for which $R(\alpha_i | x)$ is minimum

R is minimum and R in this case is called the Risk

Bayes risk = best performance that can be achieved

Two-category classification

α_1 : *deciding* ω_1

α_2 : *deciding* ω_2

$$\lambda_{ij} = \lambda(\alpha_i \mid \omega_j)$$

loss incurred for deciding ω_i when the true state of nature is ω_j

Conditional risk:

$$R(\alpha_1 \mid x) = \lambda_{11}P(\omega_1 \mid x) + \lambda_{12}P(\omega_2 \mid x)$$

$$R(\alpha_2 \mid x) = \lambda_{21}P(\omega_1 \mid x) + \lambda_{22}P(\omega_2 \mid x)$$

Minimum Risk Decision Rule

Our rule is the following:

if $R(\alpha_1 | x) < R(\alpha_2 | x)$
action α_1 : “decide ω_1 ” is taken

This results in the equivalent rule :

decide ω_1 if:

$$(\lambda_{21} - \lambda_{11}) P(x | \omega_1) P(\omega_1) > \\ (\lambda_{12} - \lambda_{22}) P(x | \omega_2) P(\omega_2)$$

and decide ω_2 otherwise

Likelihood ratio Decision Rule

The preceding rule is equivalent to the following rule:

$$\text{if } \frac{P(x|\omega_1)}{P(x|\omega_2)} > \frac{\lambda_{12} - \lambda_{22}}{\lambda_{21} - \lambda_{11}} \cdot \frac{P(\omega_2)}{P(\omega_1)}$$

Then take action α_1 (decide ω_1)

Otherwise take action α_2 (decide ω_2)

Exercise

Select the optimal decision where:

$$\Omega = \{\omega_1, \omega_2\}$$

$$\begin{aligned} P(x | \omega_1) &\longrightarrow N(2, 0.5) \text{ (Normal distribution)} \\ P(x | \omega_2) &\longrightarrow N(1.5, 0.2) \end{aligned}$$

$$P(\omega_1) = 2/3$$

$$P(\omega_2) = 1/3$$

$$\lambda = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$