

Bayes Decision Theory

Discriminant Functions for the Normal Density

Discriminant Functions for the Normal Density

- We saw that the minimum error-rate classification can be achieved by the discriminant function

$$g_i(x) = \ln P(x | \omega_i) + \ln P(\omega_i)$$

- Case of multivariate normal

$$g_i(x) = -\frac{1}{2}(x - \mu_i)^t \sum_i^{-1} (x - \mu_i) - \frac{d}{2} \ln 2\pi - \frac{1}{2} \ln |\Sigma_i| + \ln P(\omega_i)$$

Case of Independent Features with same variance

- Case 1: $\Sigma_i = \sigma^2 I$ (I stands for the identity matrix)

$g_i(x) = w_i^t x + w_{i0}$ (*linear discriminant function*)

where :

$$w_i = \frac{\mu_i}{\sigma^2}; \quad w_{i0} = -\frac{1}{2\sigma^2} \mu_i^t \mu_i + \ln P(\omega_i)$$




(ω_{i0} is called the threshold for the i th category!)

Linear Machine

- A classifier that uses linear discriminant functions is called “a linear machine”
- The decision surfaces for a linear machine are pieces of hyperplanes defined by:

$$g_i(x) = g_j(x)$$

Hyperplane Decision Boundaries



If covariance matrices for two distributions are equal and proportional to the identity matrix, then the distributions are spherical in d dimensions and the boundary is a generalized hyperplane in $d-1$ dimensions, perpendicular to the line joining the means

Hyperplane Equation

- The hyperplane separating R_i and R_j


$$x_0 = \frac{1}{2}(\mu_i + \mu_j) - \frac{\sigma^2}{\|\mu_i - \mu_j\|^2} \ln \frac{P(\omega_i)}{P(\omega_j)} (\mu_i - \mu_j)$$

always orthogonal to the line linking the means






$$\text{if } P(\omega_i) = P(\omega_j) \text{ then } x_0 = \frac{1}{2}(\mu_i + \mu_j)$$

Influence of Priors on Decision Boundaries

0.7,0.3



0.9,0.1



As priors are changed, the decision boundary shifts. For sufficiently disparate priors the boundary will not lie between the means.


- Case 2: $\Sigma_i = \Sigma$ (covariance of all classes are identical but arbitrary!)

- Hyperplane separating R_i and R_j


$$x_0 = \frac{1}{2}(\mu_i + \mu_j) - \frac{\ln[P(\omega_i)/P(\omega_j)]}{(\mu_i - \mu_j)^t \Sigma^{-1}(\mu_i - \mu_j)} \cdot (\mu_i - \mu_j)$$

(the hyperplane separating R_i and R_j is generally not orthogonal to the line between the means!)



Decision Regions for Equal but Asymmetric Gaussians



Priors
0.5, 0.5



Priors
0.9,0.1



Decision hyperplanes need not be perpendicular to line joining the means

- Case 3: Σ_i = arbitrary

- The covariance matrices are different for each category

$$g_i(x) = x^t W_i x + w_i^t x = w_{i0}$$

where :







$$W_i = -\frac{1}{2} \Sigma_i^{-1}$$

$$w_i = \Sigma_i^{-1} \mu_i$$

$$w_{i0} = -\frac{1}{2} \mu_i^t \Sigma_i^{-1} \mu_i - \frac{1}{2} \ln |\Sigma_i| + \ln P(\omega_i)$$

(Hyperquadrics which are: hyperplanes, pairs of hyperplanes, hyperspheres, hyperellipsoids, hyperparaboloids, hyperhyperboloids)

Hyperquadric Decision Boundaries



Arbitrary Gaussian distributions lead to Bayes decision boundaries that are general hyperquadrics. Conversely given any hyperquadric, one can find two Gaussian distributions whose Bayes decision boundary is that hyperquadric.
Variances are indicated by contours of constant density.