Maximum Likelihood for the HMM

Sargur Srihari srihari@cedar.buffalo.edu

HMM Topics

- 1. What is an HMM?
- 2. State-space Representation
- 3. HMM Parameters
- 4. Generative View of HMM
- 5. Determining HMM Parameters Using EM
- 6. Forward-Backward or $\alpha \beta$ algorithm
- 7. HMM Implementation Issues:
 - a) Length of Sequence
 - b) Predictive Distribution
 - c) Sum-Product Algorithm
 - d) Scaling Factors
 - e) Viterbi Algorithm

Srihari

HMM Parameters



$$p(X, Z \mid \theta) = p(z_1 \mid \pi) \left[\prod_{n=2}^{N} p(z_n \mid z_{n-1}, A) \right] \prod_{m=1}^{N} p(x_m \mid z_m, \phi)$$

where $X = \{x_1, ..., x_N\}, Z = \{z_1, ..., z_N\}, \theta = \{\pi, A, \phi\}$

We have three sets of HMM parameters: $\theta = (\pi, A, \phi)$

1.Initial Probabilities of first latent variable: Π is a vector of *K* probabilities of the states for latent variable z

2. Transition Probabilities (State-to-state for any latent variable): A is a $K \times K$ matrix of transition probabilities A_{ij}

3. Emission Probabilities (Observations conditioned on latent): ϕ are parameters of conditional distribution $p(\mathbf{x}_k | \mathbf{z}_k)$

- A and π parameters are often initialized uniformly
- Initialization of $\boldsymbol{\varphi}$ depends on form of distribution

Determining HMM Parameters

- Given data set $X = \{x_1, ..., x_n\}$ we can determine HMM parameters $\theta = \{\pi, A, \phi\}$ using maximum likelihood
- Likelihood function obtained from joint distribution by marginalizing over latent variables $Z = \{z_1, ..., z_n\}$



Computational Issues for Parameters

• Joint distribution is $p(X|\theta) = \sum_{Z} p(X, Z|\theta)$

• where

$$p(X, Z \mid \theta) = p(z_1 \mid \pi) \left[\prod_{n=2}^{N} p(z_n \mid z_{n-1}, A) \right] \prod_{m=1}^{N} p(x_m \mid z_m, \phi)$$

where $X = \{x_1, ..., x_N\}, Z = \{z_1, ..., z_N\}, \theta = \{\pi, A, \phi\}$

- Joint distribution p(X,Z|O) does not factorize over n, we cannot treat each of the summations over z_n independently
- There are *N* variables summed over each of which has *K* states, so there are *K*^N terms
 - No. of terms grows exponentially with length of chain, summing over all paths in trellis

Machine Learning

Solution to computational task

- Use conditional independence properties to reorder summations to obtain algorithm that scales linearly with length of chain
- Use Expectation Maximization to maximizing the log-likelihood function in HMMs

Srihari

EM for MLE in HMM

- 1. Start with *initial selection for model parameters* θ^{pld}
- 2. In E step take these parameter values and find posterior distribution of latent variables $p(Z|X, \theta^{old})$

Use this posterior distribution to evaluate expectation of the logarithm of the complete-data likelihood function $\ln p(X,Z \mid \theta)$

Which can be written as

$$Q(\theta, \theta^{old}) = \sum_{Z} p(Z | X, \theta^{old}) \ln p(X, Z | \theta)$$

underlined portion independent of θ is evaluated 3. In M-Step maximize *Q* w.r.t. θ

$\mbox{Expansion of } Q$

• Introduce notation Gamma and Xi

 $\gamma(z_n) = p(z_n | X, \theta^{old})$: Marginal posterior distribution of latent variable z_n

 $\xi(z_{n-1}, z_n) = p(z_{n-1}, z_n | X, \theta^{old})$: Joint posterior of two successive latent variables

• We will be re-expressing Q in terms of γ and ξ $Q(\theta, \theta^{old}) = \sum_{Z} p(Z | X, \theta^{old}) \ln p(X, Z | \theta)$



Detail of γ and ξ

For each value of *n* we can store

- $\gamma(z_n)$ using *K* non-negative numbers that sum to unity $\xi(z_{n-1}, z_n)$ using a *K* x *K* matrix whose elements also sum to unity
- Using notation

 $\gamma(z_{nk})$ denotes conditional probability of $z_{nk}^{=1}$ Similar notation for $\xi(z_{n-1,j}, z_{nk})$

 Because the expectation of a binary random variable is the probability that it takes value 1

$$\gamma(z_{nk}) = E[z_{nk}] = \sum_{z} \gamma(z) z_{nk}$$

 $\xi(z_{n-1,j}, z_{nk}) = E[z_{n-1,j}, z_{nk}] = \sum_{z} \gamma(z) \ z_{n-1,j} z_{nk}$

Expansion of Q

• We begin with

$$Q(\theta, \theta^{old}) = \sum_{Z} \underline{p(Z \mid X, \theta^{old})} \ln p(X, Z \mid \theta)$$

Substitute

$$p(X, Z \mid \theta) = p(\mathbf{z}_1 \mid \pi) \left[\prod_{n=2}^N p(\mathbf{z}_n \mid \mathbf{z}_{n-1}, A) \right] \prod_{m=1}^N p(\mathbf{x}_m \mid \mathbf{z}_m, \varphi)$$

• And use definitions of γ and ξ to get:

$$Q(\theta, \theta^{old}) = \sum_{k=1}^{K} \gamma(z_{1k}) \ln \pi_k + \sum_{n=2}^{N} \sum_{j=1}^{K} \sum_{k=1}^{K} \xi(z_{n-1,j}, z_{nk}) \ln A_{jk}$$
$$+ \sum_{n=1}^{N} \sum_{k=1}^{K} \gamma(z_{nk}) \ln p(x_n | \phi_k)$$

$$E-Step$$

$$Q(\theta, \theta^{old}) = \sum_{k=1}^{K} \gamma(z_{1k}) \ln \pi_k + \sum_{n=2}^{N} \sum_{j=1}^{K} \sum_{k=1}^{K} \xi(z_{n-1,j}, z_{nk}) \ln A_{jk}$$

$$+ \sum_{n=1}^{N} \sum_{k=1}^{K} \gamma(z_{nk}) \ln p(x_n \mid \phi_k)$$

Goal of E step is to evaluate $\gamma(z_n)$ and $\xi(z_{n-1}, z_n)$ efficiently (Forward-Backward Algorithm)



M-Step

- Maximize $Q(\theta, \theta^{old})$ with respect to parameters $\theta = \{\pi, A, \phi\}$
 - Treat $\gamma(z_n)$ and $\xi(z_{n-1}, z_n)$ as constant
- Maximization w.r.t. π and A
 - easily achieved (using Lagrangian multipliers)

$$\pi_{k} = \frac{\gamma(z_{1k})}{\sum_{j=1}^{K} \gamma(z_{1j})} \qquad A_{jk} = \frac{\sum_{n=2}^{N} \xi(z_{n-1,j}, z_{nk})}{\sum_{l=1}^{K} \sum_{n=2}^{N} \xi(z_{n-1,j}, z_{nl})}$$

- Maximization w.r.t. ϕ_k
 - Only last term of Q depends on $\phi_k \rightarrow \sum_{n=1}^{N} \sum_{k=1}^{N} \gamma(z_{nk}) \ln p(x_n | \phi_k)$
 - Same form as in mixture distribution for i.i.d.

M-Step for Gaussian Emission

- Maximization of $Q(\theta, \theta^{old})$ wrt ϕ_k
- Gaussian Emission Densities
 - $p(\mathbf{x}|\boldsymbol{\phi}_k) \sim N(\mathbf{x}|\boldsymbol{\mu}_k,\boldsymbol{\Sigma}_k)$
- Solution:



$$\Sigma_{k} = \frac{\sum_{n=1}^{N} \gamma(z_{nk}) (\mathbf{x}_{n} - \boldsymbol{\mu}_{k}) (\mathbf{x}_{n} - \boldsymbol{\mu}_{k})^{T}}{\sum_{n=1}^{N} \gamma(z_{nk})}$$

M-Step for Multinomial Observed

 Conditional Distribution of Observations have the form

$$p(\mathbf{x} \mid \mathbf{z}) = \prod_{i=1}^{D} \prod_{k=1}^{K} \mu_{ik}^{x_i z_k}$$

• M-Step equations:

$$\mu_{ik} = \frac{\sum_{n=1}^{N} \gamma(z_{nk}) x_{ni}}{\sum_{n=1}^{N} \gamma(z_{nk})}$$

 Analogous result holds for Bernoulli observed variables