

Scatterplot with large no of data points reveals little structure





Figure 3.7 A scatterplot of 96,000 cases, with much overprinting. Each data point represents an individual applicant for a loan. The vertical axis shows the age of the applicant, and the horizontal axis indicates the day on which the application was made.

Also, can conceal overprinting which can be significant for multimodal data

Contourplot overcomes some scatterplot problems



Unimodality
can be seen:
Not apparent
in scatterplot

Figure 3.9 A contour plot of the data from figure 3.7.

Requires a 2-D density estimate to be constructed
with a 2-D kernel

Display when one of the variables is time




Figure 3.10 A plot of the number of credit cards in circulation in the United Kingdom, by year.




Figure 3.12 Weight changes over time in a group of 10,000 school children in the 1930s. The steplike pattern in the data highlights a problem with the measurement process.




Figure 3.11 Patterns of change over time in the number of miles flown by UK airlines in the 1960s.




Figure 3.12 Weight changes over time in a group of 10,000 school children in the 1930s. The steplike pattern in the data highlights a problem with the measurement process.

Tools for Displaying More than Two Variables

- Scatter plots for all pairs of variables
- Trellis Plot
- Parallel Coordinates plot

More than two variables

- Sheets of Paper and Computer screens are fine for two variables
- Need projections from higher-dimensional data to 2-D plane
- Methods
 - Examine all pairs of variables
 - Scatterplot matrix
 - Trellis plot
 - Icons