

# Data Analysis and Uncertainty

## Part 3: Hypothesis Testing/Sampling

Instructor: Sargur N. Srihari

University at Buffalo  
The State University of New York  
[srihari@cedar.buffalo.edu](mailto:srihari@cedar.buffalo.edu)

# Topics

1. Hypothesis Testing
2. Sampling Methods

# Hypothesis Testing

$$\lambda = \frac{L(\theta_0 | D)}{\sup_{\varphi} L(\varphi | D)}$$

$$X^2 = \sum_{k=1,t} \frac{(E_k - O_k)^2}{E_k}$$

# Hypothesis Testing

$$p(H_i | x) \propto p(x | H_i) p(H_i)$$

$$\frac{p(H_0 | x)}{p(H_1 | x)} \propto \frac{p(H_0)}{p(H_1)} \bullet \frac{p(x | H_0)}{p(x | H_1)}$$

# Sampling Methods

$$p(\theta \mid D) \propto p(D \mid \theta)p(\theta)$$

$$p(\theta) \propto \theta^{\alpha - 1}(1-\theta)^{\beta - 1}$$

$$L(\theta \mid D) = \theta^r(1-\theta)^{n-r}$$

$$\begin{aligned}
p(\theta | D) &\propto p(D | \theta) p(\theta) \\
&= \theta^r (1 - \theta)^{n-r} \theta^{\alpha-1} (1 - \theta)^{\beta-1} \\
&= \theta^{r+\alpha-1} (1 - \theta)^{n-r+\beta-1}
\end{aligned}$$

$$\begin{aligned}
p(x(n+1) | D) &= \int p(x(n+1), \theta | D) d\theta \\
&= \int p(x(n+1) | \theta) p(\theta | D) d\theta
\end{aligned}$$

$$p(\theta | D_1, D_2) \propto p(D_2 | \theta) p(D_1 | \theta) p(\theta)$$

$$I(\theta | x) = -E\left[\frac{\partial^2 \log L(\theta | x)}{\partial \theta^2}\right]$$

$$p(\theta) \propto \sqrt{I(\theta | x)}$$

# Sampling Methods

$$\begin{aligned} & \frac{\sigma^2}{n} \left(1 - \frac{n}{N}\right) \\ & \sum (x(i) - \bar{x}) / (n - 1) \\ & \sum \frac{N_k \bar{x}_k}{N} \\ & \frac{1}{N^2} \sum N_k^2 \text{var}(\bar{x}_k) \end{aligned}$$

# Sampling Methods

$$\frac{\sum x_k / \sum n_k}{\frac{1-f}{(\sum n_k)^2} \frac{a}{1-a} (\sum s_k^2 + r^2 \sum n_k^2 - 2r \sum s_k n_k)}$$

# Means of Sample Sizes

