

# Gaussian Bayesian Networks

Sargur Srihari

[srihari@cedar.buffalo.edu](mailto:srihari@cedar.buffalo.edu)

# Topics

1. Gaussian Bayesian Network
2. Linear Gaussian Bayesian Network
3. From Gaussian distribution to Bayesian Network

# Gaussian Bayesian Networks


- We show how we can define a continuous joint distribution using a Bayesian network
  - This representation is based on the linear Gaussian model
- Definition of Gaussian Bayesian network:
  - It is a BN all of whose variables are continuous and all of the CPDs are linear Gaussians
- Surprising result:
  - linear Gaussian Bayesian networks are an alternative representation to multivariate Gaussian distributions

# Theorem of Linear Gaussian Model

- Let  $Y$  be a linear Gaussian of parents  $X_1, \dots, X_k$  :  


$$P(Y|\mathbf{x}) \sim N(\beta_0 + \beta^T \mathbf{x}; \sigma^2)$$
- Assume  $X_1, \dots, X_k$  are jointly Gaussian
  - with distribution  $N(\boldsymbol{\mu}, \Sigma)$ .
- Then distribution of  $Y$  is a normal distribution  
 $p(Y) = N(\mu_Y; \sigma_Y^2)$  where  

$$\boldsymbol{\mu}_Y = \beta_0 + \beta^T \boldsymbol{\mu}, \quad \sigma_Y^2 = \sigma^2 + \beta^T \Sigma \beta$$
- The joint distribution over  $\{ \mathbf{X}, Y \}$  is a normal distribution where  $\text{Cov}(X_i; Y) = \sum_j \beta_j \Sigma_{i,j}$ 
  - It follows that if  $\mathcal{B}$  is a linear Gaussian Bayesian network then it defines a joint Gaussian distribution<sup>4</sup>



# Example of Linear Gaussian BN

- All variables are Gaussian
- All CPDs are linear Gaussian



$$P(A) = \mathcal{N}(\mu_A, \sigma_A^2)$$

$$P(B) = \mathcal{N}(\mu_B, \sigma_B^2)$$

$$P(C) = \mathcal{N}(\mu_C, \sigma_C^2)$$


$$P(D|A, B) = \mathcal{N}(\beta_{D,0} + \beta_{D,1}A + \beta_{D,2}B, \sigma_D^2)$$

$$P(E|C, D) = \mathcal{N}(\beta_{E,0} + \beta_{E,1}C + \beta_{E,2}D, \sigma_E^2)$$

- Theorem says this defines a joint Gaussian distribution

# Ex: linear Gaussian BN to joint Gaussian

- Bayesian Network:



- CPDs:  $p(X_1) = N(1; 4)$   
 $p(X_2|X_1) = N(0.5X_1 - 3.5; 4)$   
 $p(X_3|X_2) = N(-X_2 + 1; 3)$

- From theorem, compute joint Gaussian  $p(X_1, X_2, X_3)$ :

$$\mu_2 = 0.5\mu_1 - 3.5 = 0.5 - 3.5 = -3$$

$$\mu_3 = (-1)\mu_2 + 1 = (-1)(-3) + 1 = 4$$

$$\Sigma_{22} = 4 + (1/2)^2 4 = 5$$

$$\Sigma_{33} = 3 + (-1)^2 5 = 8$$

$$\Sigma_{12} = (1/2)4 = 2$$

$$\Sigma_{23} = (-1)\Sigma_{22} = -5 \quad \Sigma_{13} = (-1)\Sigma_{12} = -2$$

$$p(X_1, X_2, X_3) = N(\mu, \Sigma) \quad \mu = \begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 4 & 2 & -2 \\ 2 & 5 & -5 \\ -2 & -5 & 8 \end{bmatrix}$$

# From joint Gaussian to a conditional

- Let  $\{X, Y\}$  have a joint normal distribution

$$p(X, Y) = N \left( \begin{pmatrix} \mu_X \\ \mu_Y \end{pmatrix}; \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \right)$$

- Then the conditional density  $P(Y|X)$  is given by

$$P(Y|X) \sim N(\beta_0 + \beta^T X; \sigma^2) \text{ where}$$

$$\begin{aligned} \beta_0 &= \mu_Y - \Sigma_{YX} \Sigma_{XX}^{-1} \mu_X \\ \beta &= \Sigma_{XX}^{-1} \Sigma_{YX} \\ \sigma^2 &= \Sigma_{YY} - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{XY} \end{aligned}$$

- Defines  $\beta$  and  $\sigma^2$  in terms of parameters of joint
  - It allows us to take a joint Gaussian distribution and produce a Bayesian network using the process of constructing an I-map

# Theorem: From Gaussian to BN


- Let  $\boldsymbol{\chi} = \{X_1, \dots, X_n\}$  and let  $p$  be a joint Gaussian distribution over  $\boldsymbol{\chi}$
- Given any ordering  $X_1, \dots, X_n$  over  $\boldsymbol{\chi}$  we can construct a Bayesian network graph  $\mathcal{G}$  and a Bayesian network  $\mathcal{B}$  over  $\mathcal{G}$  such that
  1.  $\text{Pa}_{X_i}^{\mathcal{G}} \subseteq \{X_1, \dots, X_{i-1}\}$
  2. The CPD of  $X_i$  in  $\mathcal{B}$  is a linear Gaussian of its parents
  3.  $\mathcal{G}$  is a minimal I-map for  $p$



# Ex: Gaussian to BN

- Given Gaussian
- Ordering  $X_1, X_2, X_3$
- Want to build linear Gaussian Bayes net

$$p(X_1, X_2, X_3) = N(\mu, \Sigma) \quad \mu = \begin{bmatrix} 1 \\ -4.5 \\ 8.5 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 4 & 2 & -2 \\ 2 & 5 & -5 \\ -2 & -5 & 8 \end{bmatrix}$$



- We will show that the CPDs are

$$p(X_1) = N(1 ; 4)$$

$$p(X_2 | X_1) = N(0.5X_1 - 5 ; 4)$$

$$p(X_3 | X_2) = N(-X_2 + 4 ; 3)$$


# Step 1: Determining $p(X_1)$

- Given multivariate Gaussian

$$p(X_1, X_2, X_3) = N(\mu, \Sigma) \quad \mu = \begin{bmatrix} 1 \\ -4.5 \\ 8.5 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 4 & 2 & -2 \\ 2 & 5 & -5 \\ -2 & -5 & 8 \end{bmatrix}$$

– Ordering  $X_1, X_2, X_3$

- CPD for sub-network



It can read it off from the joint Gaussian as


$$p(X_1) = N(1; 4)$$

# Step 2: Determining $p(X_2|X_1)$

- Given multivariate Gaussian

$$p(X_1, X_2, X_3) = N(\mu, \Sigma) \quad \mu = \begin{bmatrix} 1 \\ -4.5 \\ 8.5 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 4 & 2 & -2 \\ 2 & 5 & -5 \\ -2 & -5 & 8 \end{bmatrix}$$

- Ordering  $X_1, X_2, X_3$

- CPD for sub-network 

- Can read  $p(X_1, X_2)$  from joint

$$p(X_1, X_2) = N\left(\mu = \begin{bmatrix} 1 \\ -4.5 \end{bmatrix}, \Sigma = \begin{bmatrix} 4 & 2 \\ 2 & 5 \end{bmatrix}\right)$$

- But parameters for  $p(X_2|X_1)$  are computed using:

$$\begin{aligned} \beta_0 &= \mu_Y - \Sigma_{YX} \Sigma_{XX}^{-1} \mu_X \\ \beta &= \Sigma_{XX}^{-1} \Sigma_{YX} \\ \sigma^2 &= \Sigma_{YY} - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{XY} \end{aligned}$$

$$\begin{aligned} \beta_0 &= \mu_2 - \Sigma_{21} \Sigma_{11}^{-1} \mu_1 = -4.5 - (2)(1/4)(1) = -5 \\ \beta_1 &= \Sigma_{21} \Sigma_{11}^{-1} = (2)(1/4) = 1/2 \\ \sigma^2 &= \Sigma_{22} - \Sigma_{21} \Sigma_{11}^{-1} \Sigma_{12} = 5 - (2)(1/4)(2) = 4 \end{aligned}$$

- Thus  $p(X_2|X_1) = N(\beta_0 + \beta_1 X_1; \sigma^2) = N(0.5X_1 - 5; 4)$


# Step 3: Determining $p(X_3|X_2)$

- Given multivariate Gaussian

$$p(X_1, X_2, X_3) = N(\mu, \Sigma) \quad \mu = \begin{bmatrix} 1 \\ -4.5 \\ 8.5 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 4 & 2 & -2 \\ 2 & 5 & -5 \\ -2 & -5 & 8 \end{bmatrix}$$

– Ordering  $X_1, X_2, X_3$

- Desired network structure



- CPD for  $p(X_3|X_2)$

– Similar to previous step

–  $p(X_3|X_2) = N(-X_2 + 4; 3)$

# Gaussian to BN using Independencies

- Algorithm uses the principle that for  $\mathbf{Z}$  to be parents of node  $X$ ,  $(X \perp \mathbf{Y} / \mathbf{Z})$  where  $\mathbf{Y} = \mathcal{X} - \mathbf{Z}$
- Need to check if  $P(X/\mathbf{Z})$  independent of  $P(Y_i/\mathbf{Z})$ 
  - Can be done using deviance measure:
    - $\chi^2$  or mutual info (KL-div)

$$d_{\chi^2}(\mathcal{D}) = \sum_{x_i, x_j} \frac{(M[x_i, x_j] - M \cdot \hat{P}(x_i) \cdot \hat{P}(x_j))^2}{M \cdot \hat{P}(x_i) \cdot \hat{P}(x_j)}$$

$$d_I(\mathcal{D}) = \frac{1}{M} \sum_{x_i, x_j} M[x_i, x_j] \log \frac{M[x_i, x_j]}{M[x_i]M[x_j]}$$

- Simpler Method: use Information form of Gaussian

# BN from conditional independencies

Algorithm computes an I-Map which may not be unique

- $\mathcal{X} = \{X_1, \dots, X_n\}$ : set of random variables
- $\mathcal{I}$ : set of independencies

```

1  Set  $\mathcal{G}$  to an empty graph over  $\mathcal{X}$ 
2  for  $i = 1, \dots, n$ 
3     $U \leftarrow \{X_1, \dots, X_{i-1}\}$  //  $U$  is the current candidate for parents of  $X_i$ 
4    for  $U' \subseteq \{X_1, \dots, X_{i-1}\}$ 
5      if  $U' \subset U$  and  $(X_i \perp \{X_1, \dots, X_{i-1}\} - U' \mid U') \in \mathcal{I}$  then
6         $U \leftarrow U'$ 
7      // At this stage  $U$  is a minimal set satisfying  $(X_i \perp$ 
8         $\{X_1, \dots, X_{i-1}\} - U \mid U)$ 
9      // Now set  $U$  to be the parents of  $X_i$ 
10     for  $X_j \in U$ 
11       Add  $X_j \rightarrow X_i$  to  $\mathcal{G}$ 
12  return  $\mathcal{G}$ 

```

Nodes other than  $U'$