

Gaussian Markov Random Fields

Sargur Srihari

srihari@cedar.buffalo.edu

Topics

- Information Form of Gaussian
- Induced Pairwise Markov Network
 - Node and Edge potentials
- Pairwise MN as Gaussian

Representing Gaussian using undirected PGMs

- First show how a Gaussian distribution can be viewed as an MRF
- Derived almost immediately from the information form of Gaussian

Information Form of Gaussian

- **Covariance Form is:**
$$p(\mathbf{x}) = N(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}|^{1/2}} \exp\left[-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^t \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right]$$
 - Using inverse covariance matrix precision $\mathbf{J} = \boldsymbol{\Sigma}^{-1}$
 - $$-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^t \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^t \mathbf{J}(\mathbf{x} - \boldsymbol{\mu}) = -\frac{1}{2}\left[\mathbf{x}^t \mathbf{J} \mathbf{x} - 2\mathbf{x}^t \mathbf{J} \boldsymbol{\mu} + \boldsymbol{\mu}^t \mathbf{J} \boldsymbol{\mu}\right]$$
 - Since last term is constant, we get information form
$$p(\mathbf{x}) \propto \exp\left[-\frac{1}{2}\mathbf{x}^t \mathbf{J} \mathbf{x} + (\mathbf{J} \boldsymbol{\mu})^t \mathbf{x}\right]$$
- **We break-up exponent into two types of terms:**
 - Involve single variable X_i , or involve pairs X_i, X_j
 - Terms that involve only the variable X_i are:

$$\boxed{-\frac{1}{2} J_{ii} x_i^2 + h_i x_i}$$
 where the potential vector $\mathbf{h} = \mathbf{J} \boldsymbol{\mu}$
 - Terms that involve only the pair X_i, X_j are:

$$-\frac{1}{2}\left[J_{ij} x_i x_j + J_{ji} x_i x_j\right] = -J_{ji} x_i x_j$$

Induced pairwise Markov network

- Information form of Gaussian induces a pairwise Markov network
- Node potentials are derived from the potential vector h and the diagonal elements J_{ii} of the information matrix
- Edge potentials are derived from the off-diagonal entries of the information matrix
- When $J_{ij}=0$ there is no edge between X_i and X_j in the model
 - Corresponds directly to the independence assumption in the Markov network

Gaussian Markov Random Fields

- Follows directly from information form

$$p(\mathbf{x}) \propto \exp\left[-\frac{1}{2} \mathbf{x}^t J \mathbf{x} + (J \boldsymbol{\mu})^t \mathbf{x}\right]$$

– which is obtained from covariance form with $J = \Sigma^{-1}$

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left[(\mathbf{x} - \boldsymbol{\mu})^t \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})\right]$$

- Break-up exponent into two types of terms

– Using the potential vector $\mathbf{h} = J \boldsymbol{\mu}$

– Terms involving single variable X_i

$$-\frac{1}{2} J_{i,i} x_i^2 + h_i x_i$$

- Called node potentials


Terms involving pairs of variables X_i, X_j

$$-\frac{1}{2} \left[J_{i,j} x_i x_j + J_{j,i} x_j x_i \right] = -J_{i,j} x_i x_j$$

- Called edge potentials (when $J_{i,j} = 0$, there is no edge)


Converse: pairwise MN as Gaussian

- Consider any pairwise Markov network with quadratic node and edge potentials



We can write node and edge energy functions (log-potentials) as

$$\begin{aligned} \epsilon_i(x_i) &= d_0^i + d_1^i x_i + d_2^i x_i^2 \\ \epsilon_{i,j}(x_i, x_j) &= a_{00}^{i,j} + a_{01}^{i,j} x_i + a_{10}^{i,j} x_j + a_{11}^{i,j} x_i x_j + a_{02}^{i,j} x_i^2 + a_{20}^{i,j} x_j^2 \end{aligned}$$



Aggregating terms we can reformulate any set of potentials as

$$P(\mathbf{x}) = \frac{1}{Z} \exp\left(-\frac{1}{2} \mathbf{x}^T J \mathbf{x} + \mathbf{h}^T \mathbf{x}\right) \quad \text{where we assume that } J \text{ is symmetric.}$$


It defines a valid Gaussian iff J is a positive definite matrix

If so J is a legal information matrix and we can take \mathbf{h} to be a potential vector resulting in a Gaussian distribution in information matrix form

Ex: Gaussian to MN

- **Given** $p(X_1, X_2) = \mathcal{N} \left(\mu = \begin{pmatrix} 1 \\ 3 \end{pmatrix}; \Sigma = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \right)$
- **Build MN**

$$J = \Sigma^{-1} = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}$$



$$P(X_1, X_2) = \frac{1}{Z} \exp [\phi_1(X_1) + \phi_2(X_2) + \phi_{1,2}(X_1, X_2)]$$

$$\phi_1(x_1) = -\frac{1}{2} J_{1,1} x_1^2 + J_{1,1} \mu_1 x_1 = -\frac{1}{2} \cdot 1 \cdot x_1^2 + 1 \cdot 1 \cdot x_1 = -\frac{1}{2} x_1^2 + x_1$$

$$\phi_2(x_2) = -\frac{1}{2} J_{2,2} x_2^2 + J_{2,2} \mu_2 x_2 = -\frac{1}{2} \cdot 2 \cdot x_2^2 + 2 \cdot 3 \cdot x_2 = -x_2^2 + 6x_2$$

$$\phi_{1,2}(x_1, x_2) = -J_{1,2} x_1 x_2 = -(-1) \cdot x_1 x_2 = x_1 x_2$$