

Individuality of Fingerprints: Comparison of Models and Measurements

Sargur Srihari and Harish Srinivasan

TR-02-07
June 2007



University at Buffalo
The State University of New York

Center of Excellence for Document Analysis and Recognition (CEDAR)
520 Lee Entrance, Suite 202
Amherst, New York 14228

Individuality of Fingerprints: Comparison of Models and Measurements

Sargur N. Srihari and Harish Srinivasan

*Department of Computer Science and Engineering,
University at Buffalo, The State University of New York, Buffalo NY, USA
Center of Excellence for Document Analysis and Recognition (CEDAR), Buffalo NY
email:{srihari@cedar,hs32@cedar}.buffalo.edu*

Abstract

Over a hundred years, several attempts have been made to quantitatively establish the degree of individuality of fingerprints. Measurements have been made using models based on grids, ridges, fixed probabilities, relative measurements and generative distributions. This paper is a survey and assessment of various fingerprint individuality models proposed to-date. Models starting from that of Galton to recently proposed generative models are described. The models are described in terms of their attributes, similarities and differences. A detailed discussion of generative models for fingerprints, which are based on modeling the distributions of fingerprint features from a database, is given. Generative models with and without ridge information are compared. The probabilities of random correspondence arrived at by all the models are summarized. Finally, recent studies of fingerprints of twins, which strengthen the individuality argument, are discussed.

Key words: Individuality of Fingerprints, Generative models, Minutiae and Ridges, Twins Fingerprints.

1. Introduction

Fingerprints have been used for identification from the early 1900s. Their use for uniquely identifying a person has been based on two premises, that, (i) they do not change with time and (ii) they are unique for each individual. Until recently, fingerprints had been accepted by courts as a legitimate means of identification. However, after several lawsuits in United States courts, beginning with *Daubert v Merrell Dow* in 1993[1] and particularly in *USA vs Mitchell* in 1999[2], fingerprint identification has been challenged under the basis that the premises stated above have not been objectively tested and the

error rates have not been scientifically established. Though the first premise has been accepted, the second one on individuality is widely challenged.

Fingerprint individuality studies started in the late 1800s. A critical analysis of the models proposed upto about 2000 has been made by Stoney[3,4]. The goal of this paper is to provide a self-contained update on Stoney's work. This is done by providing a new organization of the models and focus on some of the newer generative models and other studies.

About twenty models have been proposed trying to establish the improbability of two random people having the same fingerprint. All of the models try to quantify the uniqueness property. Most of the models are based on minutiae. Each of these models try to find out the probability of false correspondence, i.e. probability that a wrong person is identified given a latent fingerprint collected from a crime scene from a set of previously recorded whole fingerprints, i.e., the probability that the features of two fingerprints match though they are taken from different individuals. A match here does not necessarily mean an exact match but a match within given tolerance levels.

The variety of models proposed can be classified into different categories based on the approach taken. All models establish the probability of two different people being identified as the same based on their fingerprint features– which is referred to as the probability of random correspondence (PRC). The models have been classified for better understanding based on the different approaches that have been taken through a century of individuality studies. Figure 1 shows the taxonomy, with information on which models belong to which category of models.

The models are classified into five different categories, namely, grid-based models, ridge-based models, fixed probability models, relative measurement models and generative models. Grid-based models include Galton[5] and Osterburg[6] which were proposed in the late 80s and the early 90s respectively. Ridge-based models include the Roxburgh model[7,8]. Fixed probability models contain the class of Henry-Balthazard[9,10] models. Relative measurement models include the Champod model[11] and the Trauring model[12]. In the newly introduced generative models[13–15] the distribution of fingerprint features is modeled from a database from which the PRC can then be computed.

The paper discusses models in the order of the taxonomy. Sections 2-5 discuss grid-based models, fixed probability models, ridge-based models and relative measurement models respectively. Section 6 lists features that a good fingerprint individuality model should have. Section 7 discusses generative models and also lists experiments and results obtained through implementations of such models. Section 8 contains a comparison of the PRCs derived from each of the models. Section 9 discusses the contribution of twin's studies in establishing individuality. Conclusions are given in Section 10.

2. Grid Models

Grid models use grids to divide a fingerprint into individual squares, i.e., a fingerprint is divided into squares after an enlargement step. These squares are then examined to find the distribution of minutiae. These models try to calculate the probability of occurrence of an individual square. The squares are assumed and proven to be independent of each other and therefore the probability of a particular fingerprint is calculated as the product of the probability of the occurrence of each square.




Fig. 1. Taxonomy of fingerprint individuality models based on method of analysis.

2.1. Galton's Model

Galton's[5] approach was to find the PRC by quantifying the chance of two fingerprints being from different individuals, given that the minutiae in their fingerprints are alike. Galton first tries to identify fingerprints based on the type of pattern they have, e.g., loops, whorls etc. Disagreement of the types here establish their origin from different fingers but the agreement of types only goes a short way in ascertaining their origin from the same finger. He divided fingerprints into 100 groups. While many of these could be of the same type, they have very discernible differences inbetween the groups. The fingerprints in the same group have indiscernible features.

Examining the minutiae, Galton found that a coincidence of these minutiae could be an evidence of individuality. He split the fingerprint into squares of different sizes. He assumes that the minutiae occurrence in one square is dependent on the occurrence or non-occurrence of minutiae in the neighboring squares. To avoid the complexity of non-independence, he splits a fingerprint into n-''ridge interval'' squares and tries to guess the flow of ridges in that square. The probability with which he can guess the flow of ridges correctly, given the surrounding ridges is calculated[3]. He conducted three experiments, one with tracing paper (double enlargement of fingerprint image), using a prism of the camera lucid (three-fold enlargement) and using photography and pantograph (twenty-fold enlargement). He did these experiments on 40 fingerprints with 52 trials on one or the other method mentioned above. He found that six ridge interval squares gave him a probability of 1/3 in all the three experiments. Taking into consideration some errors, he reckons this probability is to be considered. To be more accurate, the probability is 1/2 for a five-ridge interval square. That is, with five ridge interval squares, considering the surrounding ridges, there is an even chance that we can guess the flow of the ridges correctly or incorrectly. So, these squares are statistically independent.

Though some of his guesses were wrong, Galton argued that they had a very natural flow that could have happened. So, he concludes that every square also has uncertainties



Fig. 2. One of Galton's database of fingerprints. The minutiae are marked with numbers. A sample of the galton 6×6 box is shown in the figure. The area is covered and the ridge flow within the box is guessed. A 6×6 square gives an approximate probability of $1/3$ for guessing the flow correctly from the neighboring squares.

due to local incidents that the outside flow does not control. These local incidents may include enclosures or islands etc. But, it is impossible to know where they will occur. So, each square can be considered as an independent entity. There are 24 squares in each fingerprint considering 6 ridge-interval squares (5 ridge-interval squares would have been more accurate, but Galton prefers to underestimate). So, the probability of exact correspondence between two fingerprints is $(1/2)^{24}$.

To incorporate errors that can occur because of not guessing the surrounding conditions correctly, Galton included two probabilities

- Probability of not guessing correctly the general course of the surrounding ridges, b
- Probability of not guessing correctly the number of ridges that enter and exit the square, c

b is calculated to be approximately around 1/16 through the general observations in the first level classification explained above. The number of ridges that enter and exit from a square will be between 5 and 7, inclusive. Taking this into consideration, Galton guesses the probability c to be around 1/256

So, the probability of guessing the flow of fingerprints is as given in equation 1

$$\left(\frac{1}{2}\right)^{24} * \frac{1}{16} * \frac{1}{256} \tag{1}$$

The probability comes to about 1 in sixty-four thousand millions. He infers that as there are about sixteen thousand millions of humans, the probability that the fingerprint of two different persons being exactly the same is less than 1/4. When two fingerprints of each of the two persons match, then this probability becomes squared.

2.1.0.1. *Roxburgh's criticisms of Galton's Model* Roxburgh[8] criticized Galton to have been over-cautious in assuming a relation between ridges in two different squares. He argues that just because Galton sees inter-variability between the squares in a single individual, he cannot assume the same variability in different individuals. This argument could be set aside if Galton had examined enough fingerprints and saw the variability in those fingerprints as well. But, his dataset was only of size 40.

He also said that Galton overlooked other conditions that might affect the flow of ridges inside a square, apart from the surrounding conditions. He argued that even if the surrounding squares undoubtedly determined a square, they are still variable based on these other conditions. He also arrives at Galton's probabilities through an a priori (rough estimate) reasoning. Given that each 5 ridge-interval square contains one minutia, it can be guessed if it is in the outer 1/3rd of the square. So, we can guess the minutia if it is present in the outer 5/9th area. Assuming that in a 6 ridge-interval square, we have 3 minutiae in 2 squares, getting the probability to 3/8, which is approximately what Galton came up with. He argued that such rough estimates cannot be used to numerically determine the variability in the ridges. Galton's model is thus built on an assumption that the disturbance that the minutiae and ridges cause in a square leak out into the adjacent squares and thereby making them dependent variables. Roxburgh argued that this disturbance in itself will have variability that can affect the adjacent squares, like the sharpness of the bifurcation we see. This variability will increase the score of individuality. The other criticism on Galton is his inability in mentioning the degree of accuracy in what he guessed as right.

2.2. *Osterburg Model*

Osterburg's individuality model[6] also classifies fingerprints based on them being loops, whorls and arches. These classes can also be further subdivided into subclasses that will have fingerprints that appear the same to an untrained eye. Further identification involves using the Galton characteristics (minutiae). Osterburgh uses ten different minutiae types to characterize a fingerprint. They are represented in figure 2.1.

Osterburg uses 39 fingerprints to calculate the probabilities of occurrence of the above mentioned minutiae. He divides a fingerprint into 1mm x 1mm grids(as shown in figure 2.2, enlarges the image to ten times its size and counts the number and type of minutiae




Fig. 3. Representation of Galton's characteristics used by Osterburgh in his model.

in each of these grids. They sometimes also have multiple minutiae in them. The most common of them being the broken ridge which can denoted as two ending ridges. The 39 fingerprints used yielded a total of 8591 cells which could be examined. Out of these, only 23% had one or more minutiae in them. In the occupied cells, the minutiae types that were present were counted. Some cells had a combination of the above said minutiae types, for example, two dots and an ending ridge. These occurrences were rare and so can be combined as a single probability.

For his individuality model, Osterburg assumes the following:

- (i) A fingerprint is a combination of grids.
- (ii) For any cell, there are 13 possibilities: 10 minutiae, broken ridge, empty cell or any other multiple occurrence of minutiae.
- (iii) The cells are statistically independent.

From the 39 fingerprints, he calculates the probability of each of the above possible occurrences. The probability of a particular configuration is thus a multinomial distribution. If p_0 is the probability of empty cells, k_0 is the number of empty cells, p_1 is the probability of ending ridges, k_1 is the number of ending ridges..., p_{12} is the probability of multiple occurrences and k_{12} is the number of multiple occurrences, then the probability P of a given configuration is given in equation 2

$$P = p_0^{k_0} p_1^{k_1} \dots p_{12}^{k_{12}} \quad (2)$$

Each minutia type is also assigned a weight parameter that is the negative log probability for the minutia. The sum of these weights gives us the entropy of a configuration, E as in equation 3

$$E = - \sum_{i=0}^{12} k_i \log_{10} p_i \quad (3)$$

Experts frequently agree that the minimum number of minutiae needed to identify a fingerprint is 12. Osterburgh gives the benefit of doubt to the suspect by taking all these twelve minutiae as ridge endings (because it is the most common of all minutiae), the entropy of such a configuration for a 72mmsq fingerprint is nearly 20. i.e. the probability of the same configuration is 10^{-20} . The occurrence of just three trifurcations also gives us an entropy of nearly 20. Therefore, we can conclude that finding three trifurcations (rarest minutiae type) also identifies a fingerprint.




Fig. 4. Osterburg divided a print into 1mm x 1mm grids and found the occurrences of different minutiae types in these grids. Probability of a minutia type is thus calculated.

Osterburg also proves that assuming independence among the cells does not affect the overall probability. The probability of a cell having minutia increases when the surrounding cells have minutiae (from 0.26 to 0.40) but the probability increase would affect our overall entropy by one or two units. This, he argues, it is not significant as we are dealing with probabilities that are in the range 10^{-20} .

Another experiment was conducted to determine the robustness of this model with varying cell sizes. It was found that there was no big change in the probability. It is argued that, if there is an arbitrary cell size that approximates the independence relation well, then this model with cell size 1 is accurate because the probabilities do not vary with cell size.

A partial latent print obtained on a crime scene can be made my an impression of any part of a finger. Say, each template fingerprint is of size 15mm x 20mm. An input print (partial latent) of size width w mm and length l mm has $(15 - w + 1)(20 - l + 1)$ different potential places where the prints can match. Considering 10 finger prints per person, the numbers of matches are $[10(15 - w + 1)(20 - l + 1)]$. So, the decrease in entropy would be $\log_{10}[10(15 - w + 1)(20 - l + 1)]$. For a given area A , the decrease maximizes when $w = 4(A/21)^{1/2}$. For a 100mmsq area, the value comes to 2.84. For a 50 mm sq area, the value is 3.09. So, the entropy value of 20 for 12 matched ridge endings decreases to 17. For a full input fingerprint, the value decreases only by 1 (For 10 possible fingerprints).

If a is the person who committed the crime and b is a suspect and c the number of people with the same characteristics of a , then the probability of identity $P(\text{Id})$, can be denoted by $P(\text{Id}) = P(b = a | C \geq 1)$ This can be simplified to $P(\text{Id}) = E(C^{-1} | C \geq 1)$ Osterburgh takes C to be distributed hyper geometrically or binomially, with a small probability parameter. So, calculations are done assuming a poisson distribution with a small λ .

3. Fixed Probability Models

This family of models assume a fixed probability of occurrence of a minutia. The occurrence of minutia is also considered independent of each other. So, the probability of N minutia occurring at their respective places is P^N .

3.1. Henry Model

Henry[9] assumed P to be $\frac{1}{4}$ for any kind of minutia, eg: ridge ending, ridge bifurcation etc. Therefore, the probability of two finger prints to have matching minutia is $\frac{1}{4}^N$. For a fingerprint with only 10 minutiae, the probability of finding an identical fingerprint is $\frac{1}{2^2}^{10}$, i.e. one in millions.

3.2. Balthazard Model

Balthazard[10] also suggested that the probability P of a minutia occurring is $\frac{1}{4}$ based on two types of minutiae, namely **ridge ending** and **ridge bifurcation** and two directions, **left** and **right**. Each of the four possible events were assigned equal probability.

He also went ahead to calculate the number of minutiae needed to identify a person uniquely in the world population. According to his model, he concluded that 17 minutiae will be needed to identify a person conclusively in a world population of 15 billion. When the population being considered is restricted to a particular geographic location, then 11 or 12 minutiae would suffice.

3.3. Bose Model

Bose[16] also suggested the probability P to be $\frac{1}{4}$ but based on the four possibilities at each square ridge interval location, namely, **dot**, **fork**, **ending ridge** and **continuous ridge**.

3.4. Wentworth and Wilder Model

Wentworth and Wilder[17] considered four different types of minutiae, namely, ridge endings, forks, islands and breaks (A ridge ending and starting off again immediately). But, they felt that $1/4$ was a very high value for the probability of occurrence of one of these minutiae types and suggested a value of $1/50$. This was a mere guess and was not based on any experiments. Taking 9 as the number of minutiae needed to identify a

particular fingerprint (and the person it belongs to), they calculated the PRC to be one in 50⁹.

3.5. Cummins and Midlo model

Cummins and Midlo[18] adopted the Wentworth and Wilder model's P value of 1/50. They additionally bought in a "pattern factor" to account for the variation in the different fingerprint patterns. They calculated the most common fingerprint occurrence probability to be 1/31. So, the probability of two fingerprints having the same minutiae can be calculated by equation 4

$$\frac{1}{31} * \frac{1}{50}^N \tag{4}$$

3.6. Gupta model

To decide on the value of P, Gupta[19] conducted experiments with 1,000 fingerprints to find out the probability of occurrence of different types of minutiae. His experiments were aimed at finding the probability of a particular minutia at a particular position. He found that forks and ending ridges were found with a frequency of 8/100 (Approximated to 1/10) and the other minutiae were found with a frequency of 1/100. He also applied a pattern factor of 1/10 and a factor of correspondence in ridge count of 1/10.

4. Ridge-Based Models

These models use the ridge as the basis to their model. The model might go along every ridge, finding minutiae along the line and calculate the PRC from them. Though, the calculation of PRC might be similar to other models, ridge-based models start off with analysing the ridges.

4.1. Roxburgh Model

Roxburgh[7,8] draws an axis extending upward is from an origin. The axis is moved clockwise and the positions, orientation of the minutiae that are encountered are noted as shown in figure 4.1 The ridge count is also noted for each minutia. The types of minutiae noted are ridge ending and ridge bifurcation, with two possible orientations, left and right. The ridge flow is represented by assuming the ridges as (approximated) concentric circles.

Considering independence of the ridge count and type of minutiae, the number of combinations is $(RT)^n$. An additional factor P is introduced based on Galton's fingerprint classification system as the probability factor of encountering a particular fingerprint type and core type. To do away with his earlier assumption of individuality of ridge type and ridge number, he conducted extensive experiments relating the two. With this, he estimated the value of T to be 2.412 instead of 4 (2 different types with 2 orientations). So, the number of combinations is $(P)(RT)^n$. Roxburgh also considered fingerprints of poor quality. Poor quality might affect determination of minutia type and position, to




Fig. 5. Roxburgh notes the minutiae in a polar coordinate system. concentric circles one ridge apart are drawn. An imaginary axis is moved in the clockwise direction and the minutiae encountered are stored as position, orientation pairs.

overcome this, he introduced a factor Q which will take values 1-3 for decreasing quality of prints. The number of combinations become $(P)(RT/Q)^n$. An additional parameter C was introduced to account for fingerprints where the proper determination of the ridge count from the core is not possible. The factor C is the number of possible positions for the configuration. The final number of combinations is given by equation 5

$$\frac{P RT^N}{C Q} \tag{5}$$

where R is the no of concentric circles, T is the no of minutia types, N is the no of minutia, P is the estimate of probability of encountering the core type, Q is the quality factor for the fingerprint image and C is the number of possible positions for the configurations whose positions are certain.

Assuming values of the variables as $T=2.412$, $R=10$, $n=35$ (minutiae), $P=1000$, $Q=1.5$ (decent enough quality) and $C=1$ (no uncertainty about the position of configuration for the core), we have probability of duplication

$$P(\text{duplication}, 35\text{minutiae}) = 5.98 * 10^{-46}$$

To decide on the parameter n , we can consider the number of people who have access to a crime scene, look at the Probability of duplication needed and fix n as needed.

5. Relative Measurement Models

These models measure minutiae features, position and orientation, relative to other minutiae or relative to the core of the fingerprint. This helps in reducing random correspondences between minutiae which might lead to a higher PRC.

5.1. Trauring Model

Trauring's model[12] uses a system of fingerprint identification which measures the position of minutiae relative to the position of three minutiae selected while enrolling a fingerprint. He considers only two types of minutiae, ridge endings and ridge bifurcations. He assumes that they are equally probable. Their orientations, namely, left and right, are also considered equally probable. Minutiae occur at random and are independent of each other. He also assumes that two minutiae distance in relation to the reference minutia will not have a deviation of more than 1.5 ridge intervals. Given N as the number of minutiae identified, s as the minutia density (minutia per square mean pattern wavelength), r as the probability of matching reference minutiae in the false fingerprint and letting the false claimant use all the 10, the probability of false identification can be calculated as in equation 6

$$P = 10r \cdot 9\pi \frac{s}{16}^N \quad (6)$$

Trauring, through observations, calculated the value of s to be a maximum of 0.11. He also suggested a conservative number of 12 for N . r 's value was guessed to be at about 1/100. Given the values, the probability of a false identification is 4×10^{-18} .

While the individuality measurement of the model is good, it is still an identification model and requires the test minutiae to be taken using an automatic scanner. So, this model might not be helpful while considering latent prints, because of their incompleteness and low quality.

5.2. Champod and Margot model

Champod[11] designed software to search for specific minutiae in a fingerprint. One thousand good quality fingerprint images were selected for the study. Image processing algorithms were used to reduce the images into a skeletal image. A verification of the skeletal images and the original images was done to ensure the correctness of minutiae position and orientation. This was done to make sure that no connective ambiguities were introduced by the software. Nine minutiae types were considered, out of which, ridge endings and bifurcations were considered to be the primary ones. The other seven minutiae types are compound minutiae, which denote different arrangements of the two primary ones. The nine minutiae types considered were,

- (i) Ridge endings
- (ii) Ridge bifurcations
- (iii) Island, dot
- (iv) Lake
- (v) Opposed bifurcations
- (vi) Bridge

- (vii) Double bifurcations, trifurcation
- (viii) Spur
- (ix) Bifurcation opposed with an ending

The compound minutiae had a maximum distance between two basic minutiae to be considered compound

The positions of the minutiae are calculated relative to the core (using cartesian coordinates and the number of ridges between the core and that minutia). Orientation were also defined relative to other minutiae when the ridge flow was in a constant direction. The orientation is measured relative to the vertical axis.

In their analysis, they reported that density of minutiae was high in the core and delta regions. The number of minutiae was seen to follow a poisson distribution in the area above the core, but the region below the core had minor deviations from the distribution. The fingerprint was then divided into sectors of 45° with a ridge-width of five ridges. It was observed that the regions near the core had more compound minutiae than regions around the delta and the periphery. Each type's frequencies was found to be independent of the others, and the number of minutiae found. The frequencies also did not vary by finger, but mostly by types. It was observed that minutiae tend to have a direction towards the core of the fingerprint. This confirms the independence of minutiae position and orientation. The probability of a fingerprint is as in equation 7

$$P(C*) = P(N)P(T)P(S)P(L) \quad (7)$$

where N, T, S and L are number of minutiae, type, orientation and length of minutiae. Their independence hypothesis has been validated using the above mentioned experiments.

6. Stoney's Features for a Good Individuality Model

Stoney and Thornton[3,4] did a critical analysis on previous models like those of Galton, Henry-Balthazard, Osterburg, Roxburgh etc and combining them with the concerns of the FBI, came up with the features that are sought in a Fingerprint Identification Model. These features are presented below.

- (i) **Ridge Structure and Description of Minutia Location:** An individuality model should consider ridge structure to provide topological order to the fingerprint, to correct minor distortions and to provide the basis for comparing the relative position of the minutiae. Ridge structure might also be useful to provide a basis for incorporating the orientation of the minutiae in the model. Both the possible directions, namely, across the ridge flow and along the ridge flow must be considered. Also, the ridge count is invariable to quality of the fingerprint and so must be included in the model
- (ii) **Description of Minutia Distribution:** A minutia distribution is required that takes into consideration the two possible directions described above. This description of the minutia distribution should incorporate local variations in the minutia density and the variation as a result of different patterns of ridge flow. He also argues that a more fundamental relationship exists between ridge flow and minutiae. For each minutiae, that produces a ridge, there is one that consumes it. Also, imbalance in minutia orientation produces ridges that are converging or diverging. So, If the overall ridge flow is known, we also know about the distribution and

orientation of minutia because minutiae can only be present along the ridges and the orientation depends on the direction of ridge flow.

- (iii) **Orientation of Minutiae:** Minutia orientations as described above are dependent on the ridge direction. Like ridge counts they are robust to fingerprint distortion. They provide an objective criteria for comparison.
- (iv) **Variation in Minutia Type:** Minutia can be considered to be of three fundamental types, the dot, the fork and the ridge ending. Compound minutiae are also possible when minutiae occur close together. Relative frequencies of the minutiae type should be used along with the correlations between ridge flow, neighboring minutiae types and minutia density
- (v) **Variations Among Prints from the Same Source:** Fingerprints from the same finger might have variations in few features because of the errors involved in registering a fingerprint. Though the orientation and ridge count might be robust to such errors, distance between minutiae, ridge spacing and curvature of the ridge may not be. So, a criteria for tolerance of such variations should be considered. Connective ambiguities must also be allowed. Though, a connective ambiguity may not be present in all ridges in a fingerprint, it is highly probable to see a few even in the most excellent quality fingerprints. The amount of ambiguities to be tolerated should be taken as a factor based on the quality of a fingerprint image.
- (vi) **Number of Positionings and Comparisons:** The value of a fingerprint for identification is inversely proportional to the chance of false association. This depends on the number of comparisons that are attempted. The greater the number of attempts, the greater the chance for false correspondence. An attempt can include the number of fingerprints compared with and also the number of positionings possible in a single fingerprint. So, these should be included in the model as well.

7. Generative Models

Generative models are statistical models that represent the distribution of the feature. In these models, a distribution of the features is learnt through a training dataset. Features are then generated from this distribution to test their individuality. What training set is used is immaterial as long as it is representative of the entire population.

7.1. Individuality of Height

Generative models for determining individuality can be understood by considering the trivial example of using the height of a person as a biometric. The goal of the generative model for height, is to come up with an analytical value for the probability of two individuals having the same height within some tolerance $\pm\epsilon$. The steps in studying individuality using a generative model are as follows.

- (i) Consider a probabilistic generative model and estimate its parameters from a particular data set.
- (ii) Evaluate analytically the probability of two individuals to have the same height (or other bio-metric), with some tolerance $\pm\epsilon$.

For the study of individuality of height, a Gaussian density is a reasonable model, to fit the distribution of heights of individuals. Figure 6 shows modeling the heights of

individual using a Gaussian p.d.f. with mean $\mu = 5.5$ and standard deviation $\sigma = 0.5$. Now the probability of two individuals having the same height with some tolerance $\pm\epsilon$ can be derived as follows.




Fig. 6. Gaussian density used to model heights of individuals $\mu = 5.5$ and $\sigma = 0.5$ i.e. mean 5.5 feet and standard deviation 6 inches.

$$\text{Probability of one individual having height } a \pm \epsilon \text{ is } \int_{a-\epsilon}^{a+\epsilon} P(h|\mu, \sigma) dh$$

$$\text{where } P(h|\mu, \sigma) \sim \mathcal{N}(\mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

$$\text{Probability of } two \text{ individuals having height } a \pm \epsilon \text{ is } \left(\int_{a-\epsilon}^{a+\epsilon} P(h|\mu, \sigma) \right)^2$$

$$\text{Probability of two individuals having } any \text{ same height } \pm\epsilon \text{ is } \int_{-\infty}^{\infty} \left(\int_{a-\epsilon}^{a+\epsilon} P(h|\mu, \sigma) dh \right)^2 da \quad (8)$$

Equation 8 can be numerically evaluated for a given value of μ, σ . Figure 7 shows the probability values for fixed $\mu = 5.5$ (can be interpreted as 5 feet and 6 inches) and varying σ . An $\epsilon = \frac{1}{120}$ signifying a tolerance of 0.1 inches was used in the probability calculations. It is obvious to note that, when σ decreases, the width of the Gaussian is smaller and hence the probability that two individuals having the same height is more.

The PRC for height assuming a mean height of 5.5 feet and standard deviation of 0.5 inches is about 0.025. i.e. 25 out of every 1000 people have the same height (tolerance of 0.1 inches).

7.2. Generative Models for Finger Prints

Generative models for fingerprints extends on the idea of the previous section. The distributions here are much more complex. They model both the location and orientation of minutiae into a mixture model. The models are discussed in detail below.




Fig. 7. Individuality of height calculated using a Gaussian as a generative model. For different values of σ and fixed $\mu = 5.5$, the probabilities are calculated.

7.2.1. Mixture model using Hyper geometric and Binomial Distributions for minutiae

Pankanti, Prabhakar and Jain[13] assume the following while considering a fingerprint individuality model:

- Only two types of minutiae are considered, namely, ridge ending and ridge bifurcation. The two are not distinguished from each other. It is assumed that minutia orientation is neither independent of each other nor of the minutia position
- Minutiae are uniformly distributed with the restriction that they are not very close to each other.
- Correspondences between minutiae in template and input prints are independent and have an equal weight.
- Fingerprint quality has not been taken into account. Only positive matches are considered (two minutiae that match), negative matches (two minutiae don't match) are ignored
- Ridge widths are assumed to be the same
- There exists only one correct alignment between fingerprints

The data retrieved for a template and input are represented as follows

$$Template : \{\{x_1, y_1, \theta_1\}, \{x_2, y_2, \theta_2\}, \dots, \{x_m, y_m, \theta_m\}\}$$

$$Input : \{\{x'_1, y'_1, \theta'_1\}, \{x'_2, y'_2, \theta'_2\}, \dots, \{x'_n, y'_n, \theta'_n\}\}$$

Where x and y represent the position of a minutia and theta represents the orientation.

For a match between two minutia, that is, after the alignment step, the following conditions should be satisfied

$$\sqrt{(x'_i - x_j)^2 + (y'_i - y_j)^2} \leq r_0$$

$$Min(|\Theta'_i - \Theta_j|, 360 - |\Theta'_i - \Theta_j|) \leq \Theta_0$$

where r_0 is the tolerance in distance and θ_0 is the tolerance in orientation.

If A is the total area of overlap between the input and template fingerprints, and an alignment is arrived at, then if 2 minutiae are within a distance of r_0 , and have an




Fig. 8. Graphical representation of the Pankanti matching algorithm. Input print overlaps the template fingerprint. Minutiae are matched based on the location and orientation tolerance

orientation deflection of less than θ_0 then a match is reported. Graphical representation of the matching algorithm is shown in figure 7.2.1.

So, the probability of a match is given by equation 9

$$\begin{aligned}
 P(\sqrt{(x'_i - x_j)^2 + (y'_i - y_j)^2} \leq r_0) &= \frac{\text{area of tolerance}}{\text{total area}} \\
 &= \frac{\pi r_0^2}{A} = \frac{C}{A} \\
 P(\text{Min}(|\Theta'_i - \Theta_j|, 360 - |\Theta'_i - \Theta_j|) \leq \Theta_0) &= \frac{\text{angle of tolerance}}{\text{total angle}} = \frac{2\theta_0}{360}
 \end{aligned} \tag{9}$$

Therefore, the probability of only minutiae in the input matching m of the template is mC/A .

If there are two minutiae in the input, the probability of a match between one pair of minutiae and a mismatch for the other one is $\frac{mC}{A} \times \frac{A-mC}{A-C} \times 2$ (2 different possibilities of it happening, namely, first matches, second mismatch or first mismatch, second match).

If the input has n minutiae and the template has m minutiae, then the probability of matching exactly p minutiae can be given by equation 10

$$\begin{aligned}
 p(A, C, m, n, p) &= \binom{n}{p} \times \left(\frac{mC}{A}\right) \left(\frac{(m-1)C}{A-C}\right) \dots \left(\frac{(m-p+1)C}{A-(p-1)C}\right) \times \\
 &\quad \left(\frac{A-mC}{A-pC}\right) \left(\frac{A-(m-1)C}{A-(p+1)C}\right) \dots \left(\frac{(A-(m-(n-p+1))C}{A-(n-1)C}\right)
 \end{aligned} \tag{10}$$

The first p terms denote the probability of matching p minutiae and the remaining terms denote the probability of not matching the rest. Considering $M = A/C$ and assuming it to be an integer, we can reduce the above said formula to

$$P(M, m, n, p) = \frac{\binom{m}{p} \binom{M-m}{n-p}}{\binom{M}{n}}$$

This probability is only with respect to the position of minutiae. Considering orientation of the minutiae, the authors assume that it the orientation is not independent of each other for nearby minutiae, nor is it independent of the position of the minutiae. Given p minutiae have matched based on direction, the probability that q of them also match in orientation can be given by a binomial distribution (as there are only two directions possible)

$$\binom{p}{q} l^q (1-l)^{p-q}$$

where l is the probability mentioned in equation 9

So, the probability of matching q minutiae is given in equation 11

$$P(M, m, n, q) = \sum_{p=q}^{\min(m,n)} \left(\frac{\binom{m}{p} \binom{M-m}{n-p}}{\binom{M}{n}} \times \binom{p}{q} (l)^q (1-l)^{p-q} \right) \quad (11)$$

Since, minutiae can lie only on ridges, it is corrected to $M = (A/w)/2r_0$ where w is the ridge period and $2r_0$ is the length tolerance in minutia location.

In [13] the values of the parameters r_0 and θ_0 are estimated with a tolerance limit of 97.5 to be 15 pixels and 22.5° respectively. l is calculated to be 0.417 based on the value of θ_0 . w is taken to be 9.1 as reported by Stoney.

It follows the 12 point guideline used by most governments and reports the probability of false fingerprint correspondence for different values of M, m, n, q . Then calculate the probability of a correspondence when a mistake (false positive/ false negative) is committed by a fingerprint expert while matching minutiae. It is observed that the probability of false correspondence when no mistake occurs is 1.22×10^{-20} . Another important observation is that a misjudgment of a minutiae match has far more impact than missing out a minutia.

7.2.2. Mixture Model using Gaussian and Von-Mises Distributions for minutiae

Some of the shortcomings of the generative model of [13] were addressed in Dass, Zhu and Jain[14] which proposed a Gaussian mixture model to represent minutiae location and direction. Their model aims at being flexible, i.e., the model can represent the observed distributions through other fingerprint databases as well and the uncertainties can be ascertained from the model. They also assume non-independence of minutia locations and orientations. Since this is a better fit the PRC values are more reliable.




Fig. 9. Von-Mises distribution PDF

A minutia is denoted by (X,D) where X denotes the location of a minutia and D denotes the orientation. A joint distribution model for the k pairs of (X,D) can be represented by the mixture density function as in equation 12

$$f(s, \theta | \Theta_G) = \sum_{g=1}^G \tau_g f_g^X(s | \mu_g, \Sigma_g) \cdot f_g^D(\theta | \nu_g, \kappa_g, \rho_g) \quad (12)$$

where G is the number of mixture models, τ_g is the non-negative weight for that model. $f_g^X(s | \mu_g, \Sigma_g)$ is the probability density function of a bivariate Gaussian distribution over the position of minutiae. $f_g^D(\theta | \nu_g, \kappa_g, \rho_g)$ is the probability density function over the orientations and is given by

$$f_g^D(\theta | \nu_g, \kappa_g, \rho_g) = \rho_g v(\theta) \cdot I\{0 \leq \theta < \pi\} + (1 - \rho_g) v(\theta - \pi) \cdot I\{\pi \leq \theta < 2\pi\}$$

where $v(\theta)$ is the von-mises distribution modelling the angular random variables in $[0, \pi)$ with ν_g and κ_g are the von-mises parameters. Figure 7.2.2 shows the nature of Von-Mises distribution.

Each model in the mixture model represents a cluster of minutiae. The minutiae location corresponding to the g -th cluster also has its orientation correspond to the same cluster therefore establishing the interdependence between the two. Minutiae arising from the g -th component have directions that are either θ or $\theta + \pi$ and the probabilities associated with these two occurrences is ρ_g and $1 - \rho_g$ respectively. The model allows the ridge orientations to be different at different regions (different regions can be denoted by different models) while it makes sure that nearby minutiae have similar orientations (as nearby minutiae will belong to the same model).

Given a template T with n minutiae and an input/query Q with m minutiae and w out of them match, the PRC is given by

$$PRC_0 = p^*(w; Q, T) = \binom{n}{w} \cdot (p_m(Q, T))^w (1 - p_m(Q, T))^{n-w}$$

The probability is a binomial probability whose parameters are n and $p_m(Q, T)$. The latter is the probability that a random minutia from Q will match a minutia from T .

Since most of the matchers try to maximize the number of matchings (i.e. they would find a matching even in a fingerprint that are totally different, we calculate the conditional expectation, conditioned on that fact that the number of matches is always greater than zero and equating this to the number of minutia matches between Q and T, the estimation can be written as

$$\frac{n.p_m(Q, T)}{(1 - (1 - p_m(Q, T))^n)} = w_0$$

w_0 is found out by the proposed models fit into Q and T and determining the number of matches by K-plet[18] matching algorithm. Value of $p_m(Q, T)$ can be found from equation (2). There are $N(N - 1)L^2$ matches possible where the database contains N different fingers with L impressions of the same finger

$$\overline{PRC} = \frac{1}{N(N - 1)L^2} \sum_{(Q, T)_{impostor}} p^*(w; Q, T)$$

It was empirically shown in [14] that this model fits the data better than [13] in terms of all the statistical moments.

7.2.3. Generative model for minutiae and ridges

In [13,14] only minutiae were used in the framework of generative models for fingerprints. Ridge details in fingerprints provide vital information about fingerprints and it is intuitive to see that any generative model that can utilize ridge detail as well into its framework can only be a better representation of the generative model for fingerprints. Motivated by this we have formulated two possible ways to introduce ridge information to the existing generative model for minutiae discussed in section 7.2.2. A detailed discussion of these is in [15]. A fourth parameter ridge type \mathcal{T} is appended to the three existing parameters of x, y, θ of the minutiae and a generative model is built for these four parameters. The ridge type \mathcal{T} is discussed under two different ridge models. Both of these two models could be essentially summarized as: Two minutiae matching with each other on location and orientation may actually have two very different ridge shapes. Thus, to make more reliable decision on whether on earth two minutiae match with each other or not, we use ridge shape information as well as minutiae location and orientation information. The first model is a naive model using only three ridge types enabling easy learning of parameters for the generative model. The second model is a complex ridge model which uses 16 ridge types and is summarized below.

7.2.3.1. Complex Ridge Model with sixteen types The need for the complex model arose from the evidence obtained from the experiments and results[15] which do not show significant difference in the PRC values between the Generative models with and without ridge information. A more complex ridge model that includes more number of ridge types is described here. As mentioned before and in [20], a ridge is represented as ridge points, and the 6th and 12th ridge points are picked as two anchors. The reason for choosing these two numbers is described in [20]. By measuring the orientation θ of the connection from minutia to these two ridge points, they are categorized as left-backward, left-forward, right-forward and ridge-backward types[15]. We use T_i^6 and T_i^{12} to denote the RRP type of 6th and 12th ridge points on ridge R_i . T_i^6 and T_i^{12} could equals to 1, 2, 3, and 4 – the




Fig. 10. (a) ridge model - 16 types. Circles (minutiae), squares near minutiae (6^{th} RRP) and squares far from minutiae (12^{th} RRP). (b) Corresponding 16 Ridge shapes

four possible RRP types. 16 different types can be derived for a ridge and is shown in Fig. 10(a). For understanding the corresponding ridge shapes are shown in Figure 10(b). Two ridges R_i and R_j could match with each other if and only if $|T_i^6 - T_j^6| \leq 1$ and $|T_i^{12} - T_j^{12}| \leq 1$, because mostly non-linear deformation could only change an RRP type to as far as its neighbor type. We also constrain that minutiae generated by the same gaussian model have the same ridge type. A generative model can be built with a similar equation as Equ. 12 except that a new empirical distribution is now added for the ridge type parameter. The equation representing the model is given below.

$$\begin{aligned}
 & f(s, \theta, T | \Theta_G) \\
 &= \sum_{g=1}^G \tau_g f_g^X(s | \mu_g, \Sigma_g) \cdot f_g^D(\theta | \nu_g, \kappa_g, \rho_g) \cdot f_g^T(T)
 \end{aligned} \tag{13}$$

where $f_g^T(T)$ is an empirical distribution. In specific, the frequencies of ridge types in the database, will be used as the empirical distribution to replace f_g^T .

7.2.3.2. Experiments and Results Generative models without ridge information and with the complex ridge type model have been implemented and experiments have been conducted on FVC2002 DB1. The number of components G for the mixture model was found after validation using k -means clustering. The database has 100 different fingerprints with 8 impressions of the same finger. Thus, there are a total of 800 fingerprints using which the model was developed. The results are compared to that of Dass, et. al. [14]. The mixture model is fitted to the database and random fingerprints are generated from the model. Values of PRC and \overline{PRC} are calculated using the formulae listed above. The results are presented in Table 1 and 2. The PRCs are calculated through varying number of minutiae in template(m), Input(n) and the number of minutiae matched(w). Table

Table 1

PRC for different fingerprint matches with varying m (number of minutiae in template), n (number of minutiae in input) and w (number of matched minutiae) - Without ridge information. \mathbf{PRC}_0 is PRC for the general population and $\overline{\mathbf{PRC}}$ is for PRC for FVC2002-DB1.

m	n	w	\mathbf{PRC}_0	$\overline{\mathbf{PRC}}$
26	26	12	6.0×10^{-6}	7.5×10^{-5}
		16	7.8×10^{-10}	1.3×10^{-6}
		26	1.2×10^{-25}	2.8×10^{-18}
36	36	12	1.1×10^{-5}	6.0×10^{-6}
		16	3.4×10^{-9}	3.0×10^{-8}
		36	2.4×10^{-40}	9.1×10^{-24}
46	46	12	2.9×10^{-4}	1.1×10^{-5}
		16	5.7×10^{-7}	7.6×10^{-8}
		46	8.3×10^{-50}	1.7×10^{-32}

Table 2

PRC for different fingerprint matches with varying m (number of minutiae in template), n (number of minutiae in input) and w (number of matched minutiae) - With ridge information. \mathbf{PRC}_0 is PRC for the general population and $\overline{\mathbf{PRC}}$ is for PRC for FVC2002-DB1.

m	n	w	\mathbf{PRC}_0	$\overline{\mathbf{PRC}}$
26	26	12	6.0×10^{-6}	7.5×10^{-5}
		16	7.8×10^{-10}	1.3×10^{-6}
		26	1.2×10^{-25}	2.8×10^{-18}
36	36	12	3.1×10^{-8}	3.6×10^{-8}
		16	9.1×10^{-13}	2.1×10^{-11}
		36	4.2×10^{-49}	9.5×10^{-32}
46	46	12	4.3×10^{-8}	2.9×10^{-9}
		16	2.0×10^{-12}	6.6×10^{-13}
		46	1.6×10^{-67}	3.1×10^{-49}

1 shows that more the number of minutiae in the template and the input, the higher the PRC. In experiments conducted on the FVC2002 DB1, there are some differences between the results obtained here and the results in Dass, et. al. [14]. This may result from use of different matching algorithms, which w_0 depends on. It can be observed that the PRC values embedded with ridge types model are never greater than PRC values without ridge information. Table 2 shows the PRC values corresponding to use of the sophisticated ridge type model in the generative model and these probabilities are lesser when compared to those in Table 1 indicating that ridge information strengthens individuality of fingerprints.

Table 3
Comparison of different individuality models

Model	Author	Year	Sample Size	Probability of False Correspondence (N=12)	Probabilistic model	Minutiae Types
Grid	Galton	1892	75	1.45×10^{-11}	$1/16 \times 1/256 \times (1/2)^{24}$	Ending Ridge, Fork
	Osterburgh	1977	39	10^{-20}	$P = p_0^{k_0} p_1^{k_1} \dots p_{12}^{k_{12}}$	Bridge, Dot, Ending Ridge, Fork, Island, Lake, Delta, Spur, Trifurcation and Double Bifurcation
Fixed Probability	Henry	1900		5.9×10^{-8}	0.25^N	Ending Ridge, Fork
	Balthazard	1911		5.9×10^{-8}	0.25^N	Ending Ridge, Fork
	Bose	1933		5.9×10^{-8}	0.25^N	Ending Ridge, Fork, dot, continuous ridge
	Wentworth and Wilder	1918		4.1×10^{-21}	0.02^N	Ending Ridge, Fork, Island, Break
	Cummins and Midlo	1943		1.3×10^{-22}	0.032×0.02^N	
	Gupta	1968				Ending Ridge, Fork, Compound Minutiae
Ridge Based	Roxburgh	1933	80	3×10^{-16}	$(P/C)(RT/Q)^N$	Ending Ridge, Fork
Relative Measurement	Trauring	1963		4×10^{-18}	$10r(9\pi s/16)^N$	Ending Ridge, Fork
	Champod	1995	1000			Ending Ridge, Fork, Island, Lake, opposed bifurcations, Bridge, Double bifurcation, Hook, bifurcation opposed with ending
Generative	Pankanti, Prabhakar and Jain (minutiae only)	2001	2672	1.22×10^{-20}	Mixture model using Hyper geometric and Binomial distributions	Ending Ridge, Fork
	Dass, Zhu and Jain (minutiae only)	2005	2560	1.1×10^{-5}	Mixture model using Gaussian and Von-Mises distributions	Ending Ridge, Fork
	Fang, Srihari and Srinivasan (minutiae and ridges)	2006	800	3.1×10^{-8}	Mixture model using Gaussian, Von-Mises distributions and Empirical distribution for ridge type	Ending Ridge, Fork, Ridge Points

8. Comparison of different models

Different models follow different methods to calculate the PRC. Models differ in the type of minutiae to be considered. Most models consider only ridge endings and ridge bifurcations whereas some consider a wide variety of compound minutiae. This facilitates the models to get a much lower PRC. While, some models differentiate between the ridge

endings and bifurcations, some do not stating that this may lead to errors when the image quality is bad.

Models differ in the number of fingerprints they have considered for arriving at the PRC. Earlier models, because of manual analysis take very few fingerprints into consideration, thus risking the sample set not representing the general population. The newer models take more than 800 fingerprints into consideration, thus reducing this risk.

Models also differ in the approach they take in calculating the PRC. Many of the earlier models were mere guesses of probability or were very primitive. Empirical experiments, though, on very few fingerprints started with Galton. We have classified models into different categories based on their approach. Models in the same category use similar approaches, for example, both models in the generative models category use a mixture of two distributions, though the individual distributions differ.

Earlier models like Galton's divide a fingerprint into independent grids and find the PRC as a product of the probability of individual grids. Some models like Trauring's use a relative measure between minutiae for better matching. The newer models are the Generative models which try to guess the distribution of minutiae of a database and then try to find the PRC by generating fingerprints from the model. A new model utilizing ridge information incorporated into the framework of generative model has a lower PRC value.

Table 3 compares the different models based on their year of proposal, number of fingerprints used for arriving at the model, their mathematical form, types of minutiae considered and the computed PRC value.

9. Twins' Studies

The study of cohorts such as twins has been important in various physiological [21–23] and behavioral [24] settings. The genetic and environmental similarities of twins allow studies such as the effectiveness of drugs, presence of psychological traits, etc. By examining the degree to which twins are differentiated, a study may determine the extent to which a particular trait is influenced by genes or the environment. In forensics and biometrics, few twin studies have been carried out due to the lack of sufficient data. Such studies are important since any modality, e.g., fingerprints, need to be evaluated for boundary conditions where the possibility of error is maximum. Satisfactory performance with twins strengthens the reliability of the method. It also establishes the degree of individuality of the particular trait. Such an individuality measure is relevant from the viewpoint of Daubert challenges in forensic testimony [1].

Research has been done for fingerprints of twins with different data set sizes. One of earlier ones used only a single pair of twins[25]. A more recent study[26] involves a large set of fingerprints from nearly 3,000 pairs of fingers involving 206 pairs of twins[26]. The question to be answered is whether there exists a higher degree of match between individuals who are twins rather than when the individuals are not twins. The goal is to determine if friction ridge patterns from cohorts (twins) are more difficult to tell apart.

The twin study reported[26] was based on fingerprints of 297 pairs of twins. The data used in the study is of high quality and quantity due to: (i) young population considered, (ii) fingerprints were obtained using a live scan device, and (iii) there are 2,970 pairs of




Fig. 11. Distribution of level 1 features in database.

fingers. The focus of the study was on fingerprint discriminability using level 1 and level 2 features. The level 1 study was performed visually on pairs of prints. The level 2 study was based on using an automatic fingerprint identification algorithm which provided match scores in discriminating between fingerprint pairs. The level 1 results, obtained by human visual comparison, show that twins finger's have a higher probability of having the same classification (42%) than in the case of non-twins (25%). The classification refers to the ridge flow type - Arch, Tented Arch, Left loop, Right loop, Whorl and Twin loop. The distribution of level 1 feature in the database is shown in figure 11.

Level 2 features were studied using a minutiae-based matching algorithm which provides a similarity score. Distributions of scores for the same fingers of twins, and non-twins were determined. Four impostor distributions are compared: twins, non-twins, identical twins and fraternal twins. The impostor distributions (where the score corresponds to a pair of different fingers) are compared with genuine score distributions (where the pair of fingerprints are from the same finger) to study twin-discriminability. Distributions are compared using several standard statistical tests. The methodology of comparing fingerprints of twins and non-twins using Level 2 features is shown in figure 12. The statistical inferences from the level 2 study are:

- (i) The distribution of scores from a pair of twin fingers are different from those of an arbitrary pair of fingers.
- (ii) The difference between twin and non-twin score distributions is significantly smaller than that between genuine (same finger of same person taken at different times) and impostor distributions.
- (iii) The distributions of impostor scores from identical and fraternal twins are similar.

The implications of the study are that there is more similarity between twin fingers than in the case of two arbitrary fingers, that twins can be successfully discriminated using fingerprints, and that there is no significant difference between the fingerprints of identical and fraternal twins. The net result is that the argument for the individuality of fingerprints is strengthened.




Fig. 12. Methodology of comparing fingerprints of twins and non-twins using Level 2 features.

10. Summary and Conclusion

A critical analysis of several fingerprint individuality models proposed over the last century as well as very recently has been made. A taxonomy of models based on their approach to measuring individuality has been made. The top level classification of the models are: grid-based, fixed probability, ridge-based, relative measurement and generative. Early models such as those based on fixed probabilities are considered weak in argument. Recent ones, especially, the generative models are strong statistically. Generative models have been evaluated on fingerprint databases with and without ridge information. The PRC obtained for a fingerprint template and input with 36 minutiae each with 12 matching minutiae is 1.1×10^{-5} or 1 in 100,000. With the use of ridge information, this probability is smaller 3.1×10^{-8} or 1 in 33 million. The PRC is independent of the database, provided the database is representative of the whole population. It can be expected that with additional information from the fingerprint, e.g., level 3 information such as pores, the PRC of generative models can be lowered further. Finally cohort studies involving twin's fingerprints conclude that although more similar, they are discriminable, thereby strengthening the individuality argument.

11. Acknowledgement

The authors wish to thank Vinu Krishnaswamy for his valuable contribution in surveying the individuality models and Gang Fang for generating probabilities with generative models. This work was supported by the Department of Justice Grant NIJ-2005-DD-BX-K012. The opinions expressed are those of the authors and not of the DOJ.

References

- [1] U. S. Supreme Court: *Daubert v. Merrell Dow Pharmaceuticals*. (1993) US 579.

- [2] U. S. District Court Eastern District of Pennsylvania: United States of America v. Byron Mitchell. (1999) July 7, Action No. 96-407.
- [3] Stoney, D., Thornton, J.: A critical analysis of quantitative fingerprint individuality models. *Journal of Forensic Sciences* **31** (1986) 1187
- [4] Stoney, D.: Measurement of fingerprint individuality. In: *Advances in Fingerprint Technology*, 2nd Ed. By Henry C Lee, R. E Gaensslen, CRC Press (2001)
- [5] Galton, F.: *Fingerprints*. McMillan, London (1892) pp.100
- [6] Osterburg, J.: Development of a mathematical formula for the calculation of fingerprint probabilities based on individual characteristics. *Journal of American Statistical Association* **vol.772** (1997) pp.72
- [7] Roxburgh, T.: Galton's work on the evidential value of fingerprints. *Sankhya: The Indian Journal of Statistics* **vol.1** (1933) pp. 62
- [8] Roxburgh, T.: On the evidential value of fingerprints. *Sankhya: The Indian Journal of Statistics* **vol.1** (1933) pp. 89
- [9] Henry, E.: *Classification and uses of fingerprints*. Routledge & Sons London (1900) pp.54
- [10] Baltazard, V.: De l'identification par les empreintes digitales. *Comptes Rendus, des Academies des Sciences* **vol.152** (1911) pp.1862
- [11] Champod, C., Margot, P.: Computer assisted analysis of minutiae occurrences on fingerprints. *Proc. International Symposium on Fingerprint Detection and Identification*, J. Almog and E. Spinger, editors, Israel National Police, Jerusalem (1996) pp. 305
- [12] Trauring, M.: Automatic comparison of finger-ridge patterns. *Nature* (1963) 197, 938
- [13] Pankanti, S., Prabhakar, S., Jain, A.K.: On the individuality of fingerprints. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **24(8)** (2002) 10101025
- [14] Dass, S., Zhu, Y., Jain, A.K.: Statistical models for assessing the individuality of fingerprints. *Fourth IEEE Workshop on Automatic Identification Advanced Technologies* (2005) pp.3-9
- [15] Fang, G., Srihari, S., Srinivasan, H.: Generative models for fingerprint individuality using ridge types. In: *Proc. International Workshop on Computational Forensics*. (2007) Manchester, U.K.
- [16] Bose, R.: Finger prints companion. quoted in Roxburgh, T., Galton's work on the evidential value of fingerprints, *Sankhya: The Indian Journal of Statistics* **vol.1** (1933) pp.50
- [17] Wentworth, B., Wilder, H.: *Personal identification*. Richard G. Badger, Boston (1918) pp.312
- [18] Cummins, H., Midlo, C.: *Fingerprints, palms and soles*. Blakiston, Philadelphia (1943) pp.143
- [19] Gupta, S.: Statistical survey of ridge characteristics. *Int. Criminal Police Rev.* **vol.130** (1968) pp.218
- [20] Fang, G., Srihari, S.N., Srinivasan, H., Phatak, P.: Use of ridge points in partial fingerprint matching. In: *Proc. of SPIE: Biometric Technology for Human Identification IV*. (2007) pp.65390D1-65390D9
- [21] Evans, A., Caroline, G., Baal, M.V.: The genetics of coronary heart disease: the contribution of twin studies. In: *Australian Academic Press. Volume 6(5)*. (2003) 432-441
- [22] Rubocki, R., McCue, B., Duffy, K., Shepard, K., Shepherd, S., Wisecarver, J.: Natural DNA mixtures generated in fraternal twins in utero. In: *Journal of Forensic Science. Volume 46(1)*. (2001) 120-125
- [23] Splitz, E., Mountier, R., Reed, T., Busnel, M.C., Marchaland, C., Robertoux, P.L., Carlier, M.: Comparative diagnoses of twin zygosity by sslp variant analysis, questionnaire, and dermatoglyphics. *Behavioral Genetics* **26** (1996) 56-63
- [24] Goldberg, S., Perrotta, M., Minde, K., Corter, K.: Maternal behaviour and attachment in low birthweight twins and singletons. In: *Child Development. Volume 57*. (1986) 34-46
- [25] Jain, A.K., Prabhakar, S., Pankanti, S.: On the similarity of identical twin fingerprints. *Pattern Recognition* **35** (2002) 2653-2663
- [26] Srihari, S., Srinivasan, H., Fang, G.: Discriminability of fingerprints of twins. *Journal of Forensic Identification* (in press)